NRC Publications Archive Archives des publications du CNRC

Quantum oscillations in the microwave magnetoabsorption of a two-dimensional electron gas

Fedorych, O.M.; Potemski, M.; Studenikin, S.A.; Gupta, J.A.; Wasilewski, Z.R.; Dmitriev, I.A.

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

Publisher's version / Version de l'éditeur:

<https://doi.org/10.1103/PhysRevB.81.201302>

Physical Review. B, Condensed Matter and Materials Physics, 81, 20, pp. 201302-1-201302-4, 2010-05-04

NRC Publications Archive Record / Notice des Archives des publications du CNRC :

<https://nrc-publications.canada.ca/eng/view/object/?id=1c9d176b-d4c7-494a-ad24-7b1395855610>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=1c9d176b-d4c7-494a-ad24-7b1395855610>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

Questions? Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

Vous avez des questions? Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.

Quantum oscillations in the microwave magnetoabsorption of a two-dimensional electron gas

O. M. Fedorych,¹ M. Potemski,¹ S. A. Studenikin,² J. A. Gupta,² Z. R. Wasilewski,² and I. A. Dmitriev^{3,*}

¹Grenoble High Magnetic Field Laboratory, CNRS, Grenoble, France

²Institute for Microstructural Sciences, NRC, Ottawa, Ontario, Canada K1A-0R6

³Institute of Nanotechnology, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany

(Received 22 February 2010; published 4 May 2010)

We report on the experimental observation of the quantum oscillations in microwave magnetoabsorption of a high-mobility two-dimensional electron gas induced by Landau quantization. Using original resonance-cavity technique, we observe two kinds of oscillations in the magnetoabsorption originating from inter-Landau-level and intra-Landau-level transitions. The experimental observations are in full accordance with theoretical predictions. Presented theory also explains why similar quantum oscillations are not observed in transmission and reflection experiments on high-mobility structures despite of very strong effect of microwaves on the dc resistance in the same samples.

DOI: [10.1103/PhysRevB.81.201302](https://doi.org/10.1103/PhysRevB.81.201302)

PACS number(s): 73.50.Mx, 73.50.Jt, 78.20.Ls, 78.67.De

Quantum oscillations in absorption (QMA) by a two-dimensional electron gas (2DEG) in perpendicular magnetic field B , governed by the ratio ω/ω_c of the wave frequency $\omega=2\pi f_{\text{mw}}$ of external electromagnetic wave and the cyclotron frequency $\omega_c=eB/mc$, were predicted long ago by Ando¹ and observed experimentally² in the IR absorption on a low-mobility and high-density 2DEG in a Si-inversion layer. Recently, similar ω/ω_c oscillations were discovered in the dc resistance of a high-mobility 2DEG irradiated by microwaves.³ Particularly intriguing are zero-resistance states⁴ which develop in the minima of these microwave-induced resistance oscillations (MIRO).

Theoretically, both MIRO (Refs. 5–7) and QMA (Refs. 1 and 5–8) stem from microwave-assisted transitions between disorder-broadened Landau levels (LLs). However, in experiments on high-mobility samples no QMA were observed so far despite strong MIRO showing up in the same experimental conditions. Several attempts to measure microwave reflection or transmission simultaneously with MIRO reported either single cyclotron resonance (CR) peak^{9–11} or more complex structure dominated by confined magnetoplasmons (CMPs).^{12–14}

Using original resonance-cavity technique, in this work we observe well-pronounced QMA in a high-mobility GaAs/AlGaAs sample which also reveals strong MIRO in dc transport measurements. For both the T -independent QMA and dynamic Shubnikov-de Haas oscillations (SdHO), the experimental results fully agree with the theoretical predictions of Ref. 5, which generalizes theory¹ of Ando for the case of smooth disorder potential appropriate for high-mobility structures. In addition, we explain the failure to observe such quantum oscillations in transmission and reflection experiments.

We start with a summary of relevant theoretical results which includes QMA theory⁵ for dynamic conductivity at high LLs and nonlinear relation between the absorption and dynamic conductivities^{8,10,15} specific for high-mobility 2DEG samples. Consider a plane wave normally incident to the 2DEG at the interface $z=0$ between two dielectrics with permittivity ϵ_1 ($z<0$) and ϵ_2 ($z>0$). The electric field $\text{Re } \mathbf{E}_l$ of external ($l=e$), reflected ($l=r$), and transmitted ($l=t$)

waves is a real part of $\mathbf{E}_l = \mathcal{E}_l \exp(ik_l z - i\omega t) \sum_{\pm} s_{\pm}^{(l)} \mathbf{e}_{\pm}$, where the wave numbers $k_e/\sqrt{\epsilon_1} = -k_r/\sqrt{\epsilon_1} = k_t/\sqrt{\epsilon_2} = \omega/c$, coefficients $s_{\pm}^{(l)}$ describe the polarization, $\sum_{\pm} |s_{\pm}^{(l)}|^2 = 1$, and $\sqrt{2}\mathbf{e}_{\pm} = \mathbf{e}_x \pm i\mathbf{e}_y$. According to the Maxwell equations, boundary conditions at $z=0$ read $\mathbf{E}_t = \mathbf{E}_r + \mathbf{E}_e$ and $\partial_z(\mathbf{E}_t - \mathbf{E}_r - \mathbf{E}_e) = (4\pi/c^2)\partial_t \hat{\sigma} \mathbf{E}_t$. It follows that:

$$\sqrt{\epsilon_1} \mathcal{E}_e s_{\pm}^{(e)} / \mathcal{E}_t s_{\pm}^{(t)} = \sqrt{\epsilon_{\text{eff}}} + 2\pi\sigma_{\pm}/c, \quad (1)$$

where $2\sqrt{\epsilon_{\text{eff}}} = \sqrt{\epsilon_1} + \sqrt{\epsilon_2}$. Further, $\sigma_{\pm} = \sigma_{xx} \pm i\sigma_{yx}$ are the eigenvalues of the complex conductivity tensor $\hat{\sigma}$ having the symmetries $\sigma_{xx} = \sigma_{yy}$ and $\sigma_{xy} = -\sigma_{yx}$, namely, $\hat{\sigma} \mathbf{e}_{\pm} = \sigma_{\pm} \mathbf{e}_{\pm}$.

Equation (1) yields the absorption $\mathcal{A}$, transmission $\mathcal{T}$, and reflection $\mathcal{R}$ coefficients (see also Refs. 8, 10, and 15),

$$\mathcal{A} = \sum_{\pm} \frac{\sqrt{\epsilon_1} |s_{\pm}^{(e)}|^2}{|\sqrt{\epsilon_{\text{eff}}} + 2\pi\sigma_{\pm}/c|^2} \text{Re} \frac{4\pi\sigma_{\pm}}{c}, \quad (2)$$

$$\mathcal{T} = \sum_{\pm} \frac{\sqrt{\epsilon_1 \epsilon_2} |s_{\pm}^{(e)}|^2}{|\sqrt{\epsilon_{\text{eff}}} + 2\pi\sigma_{\pm}/c|^2}, \quad (3)$$

$$\mathcal{R} = \sum_{\pm} |s_{\pm}^{(e)}|^2 \left| \frac{\sqrt{\epsilon_1} - \sqrt{\epsilon_2} - 4\pi\sigma_{\pm}/c}{\sqrt{\epsilon_1} + \sqrt{\epsilon_2} + 4\pi\sigma_{\pm}/c} \right|^2. \quad (4)$$

It is important to mention that the dynamic conductivity $\sigma_{\pm}$, which is the focus of present study, is the response to the screened electric field acting on 2D electrons. By contrast, coefficients in Eqs. (2)–(4) represent a response to the (unscreened) electric component of incoming wave and, therefore, measure both single-particle (transport) and collective (screening) properties of 2DEG.

The dynamical screening, represented by the denominators in Eqs. (2)–(4), becomes particularly strong in high-mobility structures where the ratio $|2\pi\sigma_{\pm}/c|$ reaches values much larger than unity. Indeed, in the absence of Landau quantization the conductivity $\sigma_{\pm} = \sigma_{\pm}^D$ is given by the Drude formula

$$\sigma_{\pm}^D = \frac{ne^2/m}{\tau_{tr}^{-1} - i(\omega \pm \omega_c)}, \quad (5)$$

where τ_{tr} is the momentum relaxation time. The absorption, Eq. (2), takes the form

$$A^D = \sqrt{\frac{\epsilon_1}{\epsilon_{eff}}} \sum_{\pm} |s_{\pm}^{(e)}|^2 \frac{\Omega \tau_{tr}^{-1}}{(\Omega + \tau_{tr}^{-1})^2 + (\omega \pm \omega_c)^2}, \quad (6)$$

where $\hbar\Omega = 2\alpha\epsilon_F/\sqrt{\epsilon_{eff}}$, $\alpha = e^2/\hbar c \simeq 1/137$ is the fine-structure constant, and ϵ_F is the Fermi energy of 2DEG. In high-mobility 2DEG $\Omega\tau_{tr} \gg 1$ and the width of the cyclotron peak in Eqs. (2)–(4) and (6) is dominated by strong reflection of microwaves. In the region $|\omega - \omega_c| \lesssim \Omega$, where $|2\pi\sigma_{\pm}| \gg c$, the collective effects are pronounced. In this region, a special care should be taken to avoid the finite-size magnetoplasmon effects.^{12–14}

According to Ref. 5 (which generalizes the results of Ref. 1 for the relevant case of smooth disorder potential, see also Refs. 7 and 8), Landau quantization at high LLs modifies Drude formula to the form which we call the quantum Drude formula (QDF) in what follows:

$$\text{Re } \sigma_{\pm} = \frac{ne^2}{\omega m} \int \frac{d\varepsilon (f_{\varepsilon} - f_{\varepsilon+\omega}) \tilde{\nu}(\varepsilon) \tau_{tr,B}^{-1}(\varepsilon + \omega)}{[\tau_{tr,B}^{-2}(\varepsilon) + \tau_{tr,B}^{-2}(\varepsilon + \omega)]/2 + (\omega \pm \omega_c)^2}, \quad (7)$$

where f_{ε} is the Fermi distribution function. The Landau quantization leads to the oscillatory density of states (DOS), $\nu(\varepsilon) = \nu(\varepsilon + \omega_c) \equiv \nu_0 \tilde{\nu}(\varepsilon)$, and to renormalization of the transport relaxation time, $\tau_{tr,B}(\varepsilon) \equiv \tau_{tr}/\tilde{\nu}(\varepsilon)$, where $\nu_0 = m/2\pi\hbar^2$ is the zero- B DOS per spin orientation. In the limit of strongly overlapping LLs, $\omega_c\tau_q \ll 1$, where τ_q is the quantum relaxation time, the DOS is weakly modulated by magnetic field

$$\tilde{\nu}(\varepsilon) = 1 - 2\delta \cos \frac{2\pi\varepsilon}{\omega_c}, \quad \delta = e^{-\pi/\omega_c\tau_q} \ll 1. \quad (8)$$

In the opposite limit $\omega_c\tau_q \gg 1$, LLs become separated, $\tilde{\nu}(\varepsilon) = \tau_q \text{Re} \sqrt{\Gamma^2 - (\varepsilon - \varepsilon_n)^2}$, where $\Gamma = \sqrt{2\omega_c/\pi\tau_q} < \omega_c/2$, and ε_n marks the position of the nearest LL.

At the CR $\omega = \omega_c$, QDF [Eq. (7)] reads

$$\sigma_{-}|_{\omega=\omega_c} = \sigma_{-}^D|_{\omega=\omega_c} \int d\varepsilon \Theta[\nu(\varepsilon)] \frac{f_{\varepsilon} - f_{\varepsilon+\omega}}{\omega}, \quad (9)$$

where the integration is over the regions with $\nu(\varepsilon) > 0$. In the case of overlapping LLs, this produces the classical Drude result $\sigma_{-} = \sigma_{-}^D$ meaning that quantum effects in the vicinity of the resonance are absent. In the case of separated LLs, $\omega_c \gg \Gamma$, the conductivity is reduced, $\sigma_{-}|_{\omega=\omega_c} = (2\Gamma/\omega_c)\sigma_{-}^D$. At the same time, the CR width increases from τ_{tr}^{-1} to $\tau_{tr}^{-1}\omega_c/\Gamma$.^{5,16}

In what follows we consider the region $|\omega - \omega_c| \gg \tau_{tr}^{-1}$ where one can safely neglect τ_{tr}^{-1} in the denominator of Eq. (6) and QDF, Eq. (7), which gives

FIG. 1. (Color online) Magnetoabsorption calculated using Eqs. (6) and (10) for two temperatures $T=2$ and 0.5 K. Frequency $f_{mw}=60$ GHz, density $n=1.8 \times 10^{11} \text{ cm}^{-2}$, mobility $\mu=10^7 \text{ cm}^2/\text{V s}$, and $\tau_q=15$ ps. The difference between traces at $T=0.5$ and 2 K in the inset shows the T -dependent dynamic SdHO.

$$\mathcal{A}/\mathcal{A}^D = \int d\varepsilon \frac{f_{\varepsilon} - f_{\varepsilon+\omega}}{\omega} \tilde{\nu}(\varepsilon) \tilde{\nu}(\varepsilon + \omega). \quad (10)$$

Equation (10) is the key theoretical result for our study. It expresses the imbalance between the rates of absorption and emission of the microwave quanta, both proportional to the product of initial and final densities of states. The screening properties of a 2DEG at $|\omega \pm \omega_c| \gg \tau_{tr}^{-1}$ are solely determined by the nondissipative part of conductivity, $\text{Im } \sigma_{\pm} \simeq ne^2/m(\omega \pm \omega_c) \gg \text{Re } \sigma_{\pm}$, which remains finite at $\tau_{tr}^{-1} \rightarrow 0$. That is why the transmission and reflection coefficients, Eqs. (3) and (4), in high-mobility structures are almost independent of disorder scattering making it very difficult to observe QMA in direct transmission and reflection experiments.^{9–13}

Magnetoabsorption at two temperatures $T=2$ and 0.5 K as given by Eqs. (6) and (10) is illustrated in Fig. 1 for high-mobility 2DEG similar to best structures used for MIRO measurements. The broadening of the envelope Drude peak [Eq. (6)] is dominated by strong reflection of microwaves near the CR since in our example $\Omega \sim \omega \gg \tau_{tr}^{-1}$. The Drude peak in Fig. 1 is modulated by the dynamic SdHO (at $\omega_c \gtrsim \omega$ and $T=0.5$ K) and by the temperature-independent QMA with maxima at the CR harmonics $\omega = N\omega_c$, which we discuss below.

At low temperature, $X_T = 2\pi^2 T/\hbar\omega_c \lesssim 1$, the absorption [Eq. (10)] manifests the dynamic SdHO, which in the case of overlapping LLs [Eq. (8)] are given by

$$\mathcal{A}/\mathcal{A}^D = 1 - \frac{4X_T\delta}{\sinh X_T} \frac{\omega_c}{2\pi\omega} \sin \frac{2\pi\omega}{\omega_c} \cos \frac{2\pi\varepsilon}{\omega_c}. \quad (11)$$

The dynamic SdHO are exponentially suppressed at $X_T \gg 1$ similar to SdHO in the dc resistance. In the inset it is clearly seen that the dynamic SdHO are periodically modulated according to $\sin(2\pi\omega/\omega_c)$ with nodes at the CR harmonics.

At $e^{-X_T} \ll \delta$, i.e., $T \gg T_D = \hbar/2\pi\tau_q$, SdHO [Eq. (11)] become exponentially smaller than T -independent ω/ω_c oscillations of second order $\mathcal{O}(\delta^2)$ which represent QMA,

FIG. 2. (Color online) MRO measured on sample 1 for $f_{\text{mw}}=67$ GHz at $T=2$ K (inset), and the amplitude of MRO vs ω/ω_c in semilog scale (main panel). The linear fit yields the quantum-scattering time $\tau_q=9.1$ ps, see Eqs. (8) and (13).

$$\mathcal{A}/\mathcal{A}^D \approx \langle \tilde{\nu}(\varepsilon) \tilde{\nu}(\varepsilon + \omega) \rangle = 2\delta^2 \cos \frac{2\pi\omega}{\omega_c}. \quad (12)$$

Here the angular brackets denote ε —averaging over the period ω_c . Maxima of QMA seen in Fig. 1 appear at integer harmonics of the CR $\omega/\omega_c=N$. The amplitude of QMA [Eq. (12)] becomes of order unity when the DOS modulation is pronounced, i.e., $\delta \sim 1$.¹⁷

Since QMA lie in the microscopic origin of MRO,⁵ it is instructive to compare the expression (12) to results for MRO in the same regime. For inelastic mechanism of MRO,^{5,6} the photoresistivity ρ_{ph} in terms of Drude resistivity ρ^D reads

$$\frac{\rho_{\text{ph}}}{\rho^D} = 1 + 2\delta^2 - \frac{4\tau_{\text{in}}\mathcal{P}^D}{\omega^2\nu_0} \delta^2 \frac{2\pi\omega}{\omega_c} \sin \frac{2\pi\omega}{\omega_c}, \quad (13)$$

where the microwave power dissipated in the absence of Landau quantization is $\mathcal{P}^D = \mathcal{A}^D \sqrt{\epsilon_1 c E_e^2} / 4\pi$ and τ_{in} is the inelastic relaxation time. Below we use this equation to determine τ_q from dc measurement of MRO. The known value of τ_q enables a direct comparison of the measured and calculated QMA [Eq. (12)] without fitting parameters.

Experiment. Two rectangular samples were cleaved from a single molecular beam epitaxy-grown wafer of a high-mobility GaAs/AlGaAs heterostructure V0050. After illumination with red light the electron concentration was $n=3.6 \times 10^{11} \text{ cm}^{-2}$ and mobility $\mu=5 \times 10^6 \text{ cm}^2/\text{V s}$. Sample 1 was prepared for dc transport measurements. In-Sn Ohmic contacts were made by rapid annealing in reducing atmosphere of argon bubbled through hydrochloric acid. The sample was placed in a helium cryostat equipped with a superconducting magnet. Similar to Refs. 10 and 12, microwaves from an HP source, model E8257D, were delivered to the sample using Cu-Be coaxial cable terminated with a 3 mm antenna.

Inset in Fig. 2 shows dc resistance R_{xx} measured in sample 1 at $f_{\text{mw}}=70$ GHz and $T=2$ K which displays well-resolved MRO. Main panel in Fig. 2 presents the dependence of $\log_{10}(\Delta R_{xx} \omega_c / \omega)$ vs ω/ω_c , where ΔR_{xx} is the amplitude of MRO measured between adjacent peaks and dips.

FIG. 3. (Color online) Magnetoabsorption measured on sample 2 at $T=2$ K for $f_{\text{mw}}=58.44$ GHz (top curve) and absorption coefficient $\mathcal{A}$ [Eqs. (6) and (10), bottom curve] calculated without fitting parameters using $\tau_q=9.1$ ps determined from MRO measurements on sample 1 (Fig. 2).

In accordance with Eq. (13) containing $\delta^2 = \exp(-2\pi/\omega_c \tau_q)$, this dependence is linear. The slope gives the quantum relaxation time $\tau_q=9.1$ ps which we use for calculation of QMA in sample 2.

Sample 2 was cleaved from the same wafer in vicinity to sample 1. In order to reduce the CMP effects,^{14,15,18} which can obscure weak QMA oscillations under the investigation, we cleaved a narrow $0.5 \times 1.5 \text{ mm}^2$ rectangular stripe. The magnetoabsorption experiment was performed using a home-built microwave cavity setup at liquid-helium temperatures.¹⁹ The cavity with a tunable resonance frequency had a cylindrical shape with 8 mm diameter and the height between 3 and 8 mm adjustable with a movable plunger. The cavity operated in TE_{011} mode, where $\{011\}$ are the numbers of half-cycle variations in the angular, radial, and longitudinal directions, respectively. The 2DEG stripe was placed at the bottom of the cavity with the external magnetic field normal to the 2DEG plane and the microwave electric field of the TE_{011} mode oriented along the short side of the rectangular sample. The sample was placed in a “face up” fashion such that the active 2DEG layer is separated from the plunger surface by the substrate. This geometry has higher sensitivity to weak-absorption signals such as QMA but is not appropriate to study, i.e., CMPs in the vicinity of the CR due to cavity over-coupling effects. To further improve sensitivity we measure the differential signal with respect to magnetic field.

The top curve in Fig. 3 presents B dependence of the second derivative of the measured absorption for $f_{\text{mw}}=58.44$ GHz whereas the bottom curve shows the second B derivative of the absorption coefficient $\mathcal{A}$, Eqs. (6) and (10), calculated without fitting parameters using $\tau_q=9.1$ ps determined from MRO measurements on sample 1 (Fig. 2). Both curves demonstrate well-resolved QMA with maxima at harmonics of the CR and dynamic SdHO with maxima determined by the position of the chemical potential with respect to LLs. Clearly, the theory reproduces the experimental trace in Fig. 3 quite well everywhere except for the shaded region around the CR where the signal is distorted due to the CMP absorption. The dimensions of sample 2

FIG. 4. (Color online) QMA measured at different frequencies compared to the theory [Eq. (12), parameters as in Fig. 3].

were chosen to allow only one CMP mode at $B=0.092$ T. The absence of higher modes for $f_{\text{mw}} < 63$ GHz enabled the possibility to observe clear quantum oscillations on both sides of the shaded region where finite-size effects^{14,15,18} are not essential.

The low-field traces ($\omega_c/\omega < 1/2$) of the magnetoabsorption are shown in Fig. 4 for several microwave frequencies together with the function $\delta^2 \cos(2\pi\omega/\omega_c)$. The phase and B damping of the observed QMA follow well the theoretical

dependence, Eq. (12), without fitting parameters. We, therefore, believe that the observed oscillations are indeed QMA predicted in Refs. 1 and 5, which provides an important experimental evidence supporting the theory of MIRO (Refs. 5–7) based on inter-LL transitions. In our sample, QMA are strongly damped [$\delta^2=0.02$ at $\omega_c=\omega/2$, $\omega/2\pi=f_{\text{mw}}=58.44$ GHz, and $\tau_q=9.1$ ps, see Eq. (12)] which makes their observation difficult. We expect that much stronger QMA as well as dynamic SdHO can be observed on samples with higher mobility (longer τ_q), as simulated in Fig. 1, provided the CMP effects are avoided or sufficiently reduced.

In summary, we have observed quantum magneto-oscillations in the microwave absorption and dynamic SdHO in a high-mobility 2DEG. For this purpose we used a sensitive high- Q cavity technique and developed a special setup to avoid undesirable magnetoplasmon effects masking the quantum oscillations. Using the quantum Drude formula⁵ and the quantum relaxation time extracted from the MIRO measurements on the same wafer we were able to reproduce the experimental results for absorption without fitting parameters, which provides a strong experimental support to the theory of MIRO and QMA based on inter-LL transitions.

We are thankful to A. D. Mirlin, D. G. Polyakov, B. I. Shklovskii, S. A. Vitkalov, and M. A. Zudov for fruitful discussions. This work was supported by the DFG, by the DFG-CFN, by Rosnauka under Grant No. 02.740.11.5072, by the RFBR, and by the NRC-CNRS project.

*Also at Ioffe Physical Technical Institute, 194021 St. Petersburg, Russia.

¹T. Ando, *J. Phys. Soc. Jpn.* **38**, 989 (1975).

²G. Abstreiter, J. P. Kotthaus, J. F. Koch, and G. Dorda, *Phys. Rev. B* **14**, 2480 (1976).

³M. A. Zudov, R. R. Du, J. A. Simmons, and J. L. Reno, *Phys. Rev. B* **64**, 201311(R) (2001); P. D. Ye *et al.*, *Appl. Phys. Lett.* **79**, 2193 (2001).

⁴R. G. Mani *et al.*, *Nature (London)* **420**, 646 (2002); M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **90**, 046807 (2003).

⁵I. A. Dmitriev, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. Lett.* **91**, 226802 (2003).

⁶I. A. Dmitriev, M. G. Vavilov, I. L. Aleiner, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. B* **71**, 115316 (2005).

⁷M. G. Vavilov and I. L. Aleiner, *Phys. Rev. B* **69**, 035303 (2004).

⁸O. E. Raichev, *Phys. Rev. B* **78**, 125304 (2008).

⁹J. H. Smet, B. Gorshunov, C. Jiang, L. Pfeiffer, K. West, V. Umansky, M. Dressel, R. Meisels, F. Kuchar, and K. von Klitzing, *Phys. Rev. Lett.* **95**, 116804 (2005).

¹⁰S. A. Studenikin, M. Potemski, A. Sachrajda, M. Hilke, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **71**, 245313 (2005).

¹¹L. C. Tung *et al.*, *Solid State Commun.* **149**, 1531 (2009).

¹²S. A. Studenikin *et al.*, *IEEE Trans. Nanotechnol.* **4**, 124 (2005).

¹³A. Wirthmann, B. D. McCombe, D. Heitmann, S. Holland, K. J. Friedland, and C. M. Hu, *Phys. Rev. B* **76**, 195315 (2007).

¹⁴S. A. Studenikin, B. D. McCombe, D. Heitman, S. Holland, K. J. Friedland, and C. M. Hu, *Phys. Rev. B* **76**, 165321 (2007).

¹⁵K. W. Chiu, B. D. McCombe, D. Heitmann, S. Holland, K. J. Friedland, and C. M. Hu, *Surf. Sci.* **58**, 182 (1976).

¹⁶M. M. Fogler and B. I. Shklovskii, *Phys. Rev. Lett.* **80**, 4749 (1998).

¹⁷Resonant features at overtones of CR can be induced by the quasiclassical memory effects even in the absence of Landau quantization, $\delta \rightarrow 0$ [see I. A. Dmitriev, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. B* **70**, 165305 (2004)]. These classical oscillations originate from the non-Markovian memory effects in the presence of both long-range and short-range randomness in the static impurity potential. The predicted phase of classical oscillations in the dynamic conductivity (and therefore in the absorption coefficient) is opposite to the phase of quantum oscillations [Eq. (12)] while the amplitude can be comparable to the value of Drude conductivity.

¹⁸O. M. Fedorych *et al.*, *Int. J. Mod. Phys. B* **23**, 2698 (2009).

¹⁹M. Seck and P. Wyder, *Rev. Sci. Instrum.* **69**, 1817 (1998).

American Physical Society

[Log in](#) | [Create Account](#) ([what's this?](#))

[RSS Feeds](#) | [Email Alerts](#)

[Home](#) [About](#) [Current Issue](#) [Archives](#) [For Contributors](#) [APS Journals](#)

SEARCH

Coming Soon in *Physics*

- Waveguides: Optical analog for a Luttinger liquid
- Materials science: Grain boundary growth

Now in Focus

[How to Shout Without Anyone Hearing](#)

April 1, 2011

Theorists can mathematically construct waves that propagate on straight-line paths through environments with many obstacles that would ordinarily cause scattering.

Feedback

Let us know what you think of *Physics*. Please email physics@aps.org with your comments, ideas, or suggestions for topics.

[APS](#) » [Journals](#) » [Physics](#) » [Synopsis](#)es » Quantum oscillations in a high-mobility electron gas

Quantum oscillations in a high-mobility electron gas

[Quantum oscillations in the microwave magnetoabsorption of a two-dimensional electron gas](#)

O. M. Fedorych, M. Potemski, S. A. Studenikin, J. A. Gupta, Z. R. Wasilewski,
and I. A. Dmitriev

Phys. Rev. B **81**, 201302 (Published May 4, 2010)

[Semiconductor Physics Mesoscopics Nanophysics](#)

Illustration: Courtesy of O. Fedorych *et al.*

Over the past decade, many nonequilibrium phenomena have been discovered in high-mobility quantum Hall systems in a perpendicular magnetic field. For instance, microwave illumination of a two-dimensional electron gas leads to oscillations of the dissipative resistance with magnetic field (MIRO). When the microwave power is sufficiently strong, the net resistance in the minima of MIRO becomes vanishingly small, which has been attributed to electrical instability of a homogeneous state with negative absolute resistivity leading to a domain structure with zero net resistance.

MIRO originate from microwave-induced transitions between quantized Landau levels broadened by disorder. Such transitions should also lead to quantum magneto-oscillations in the absorption (QMA) of electromagnetic waves, as predicted more than thirty years ago by Ando and observed in low-mobility Si structures. Nevertheless, so far all attempts to detect QMA in high-mobility systems have been unsuccessful, despite the fact that dc transport measurements in the same samples revealed pronounced MIRO. The difficulty arises from the large metallic screening of microwaves by a high-mobility two-dimensional electron gas and also from absorption at magnetoplasmon resonances that mask QMA.

Now, in a paper appearing as a Rapid Communication in *Physical Review B*, Oleh Fedorych and colleagues in Grenoble, France, working with collaborators in Canada and Germany, have succeeded in simultaneous observation of QMA and MIRO in the same high-mobility quantum Hall system using a sensitive high-Q cavity technique. The observations are in a perfect agreement with a microscopic theoretical model that relates both types of oscillations to electron kinetics involving microwave-assisted inter- and intra-Landau level transitions. – *Brad Rubin*

[Contact Us](#) | [FAQ](#) | [Privacy](#) | [Policies](#) | [For Contributors](#) | [APS Journals](#) | [RSS Feeds](#) | [APS](#)

ISSN 1943-2879. Use of the [American Physical Society](#) websites and journals implies that the user has read and agrees to our [Terms and Conditions](#) and any applicable [Subscription Agreement](#).