NRC Publications Archive Archives des publications du CNRC

Solution to the Avogadro constant challenge

Jensen, William B.; Meija, Juris

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

Publisher's version / Version de l'éditeur:

<https://doi.org/10.1007/s00216-010-3961-8>

Analytical and Bioanalytical Chemistry, 398, 1, pp. 11-12, 2010-07-29

NRC Publications Record / Notice d'Archives des publications de CNRC:

<https://nrc-publications.canada.ca/eng/view/object/?id=3fc37ab8-1a94-4d0a-9efb-a54803858098>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=3fc37ab8-1a94-4d0a-9efb-a54803858098>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

Questions? Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

Vous avez des questions? Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.

Analytical Challenge

Solution to the Avogadro constant challenge

William B. Jensen¹ and Juris Meija²

(1) Department of Chemistry, University of Cincinnati, Cincinnati, OH 45221, USA

(2) Institute for National Measurement Standards, National Research Council Canada, 1200 Montreal Road, Ottawa, ON, K1A 0R6, Canada

Email: juris.meija@nrc.ca

The Avogadro constant has long been defined as the number of molecules of a substance in a gram molecular weight [[1](#)]. In a modern parlance, it is a fundamental physical constant representing the number of entities comprising 1 mol. It is clear that any modification to the definition of macroscopic (kilogram) or microscopic mass scales (atomic mass) will affect the numerical value of the Avogadro constant. The definition of kilogram has remained unaltered since the very first *Conférence Générale des Poids et Mesures* in September of 1889, when it was declared that the international prototype of the kilogram “shall henceforth be considered to be the unit of mass,” not so with the definition of the atomic mass scale. In 1803/1805, John Dalton established the first atomic mass scale in which hydrogen was assigned $A_r(\text{H}) = 1$, whereas the current definition sets $A_r(^{12}\text{C}) = 12$. In addition, there were *numerous* other scales used during the intervening years (Table [1](#)). Though Dalton's $\text{H} = 1$ mass scale was used in chemistry for nearly a century, many of these alternative scales, following a suggestion by Wollaston, employed oxygen as the standard instead—an idea which resurfaced near the end of the nineteenth century as the $\text{O} = 16$ scale. With the discovery of isotopes, the oxygen scale was refined to the ^{16}O isotope, albeit only among physicists.

Table 1 Relative atomic mass scales through the centuries

19th century: average mass scale

H = 1 (O = 5.5)	Dalton, 1803–5
H = 1 (O = 16)	Davy, 1812
O = 10	Wollaston, 1813
O = 1	Thomson, 1813, 1825
O = 100	Berzelius, 1814
O = 4	Griffin, 1834
O = 16	Clarke, 1893
C = 12	Hinrichs, 1893
20 th century: isotopic mass scale	
¹⁶ O = 16	Aston, 1931
¹² C = 12	IUPAC, 1961

Notations O = 16 or ¹²C = 12 describe the relative atomic mass (atomic weight) scale. When $m(E)$ is the average mass of atoms of element E, the atomic weight of E is given by $A_r(E) = m(E)/m_u$, where m_u is the atomic mass constant. In the O = 16 scale $m_u = m(O)/16$, whereas $m_u = m(^{12}C)/12$ in the ¹²C = 12 scale

The most recent change in the atomic mass scale occurred during the height of the cold war when the oxygen scale, $A_r(^{16}O) = 16$, was abandoned in favor of the current $A_r(^{12}C) = 12$ by the International Union of Pure and Applied Physics (Ottawa 1960) and by the International Union of Pure and Applied Chemistry (Montreal 1961) [2].

The relative atomic mass of the ¹⁶O isotope in the ¹²C-scale is 15.9949—a difference of 0.03% from the “old” value of 16.0000. This shift, in turn, alters the value of the Avogadro constant from 6.024×10^{23} to $6.022 \times 10^{23} \text{ mol}^{-1}$. This “change” in the numerical value of the Avogadro constant can be easily spotted by inspection of twentieth century chemistry textbooks [3].

It may be of interest here to mention an earlier story regarding the value of the Avogadro constant when determination of N_A involved measurements of lattice spacing in crystals [4]. In 1919, at a time when wavelengths could not be measured directly, Manne Siegbahn (1924 Nobel Prize in Physics) proposed the 'local' non-SI unit of length for measuring the wavelength of x-rays. A student of Siegbahn, Erik Bäcklin, in his 1928 dissertation showed that wavelengths measured at incidence angles near grazing were 0.2% higher than the equivalent wavelengths measured by traditional crystal diffraction methods. This single discrepancy initially challenged the work of no less than five Nobel laureates—Richards, Bragg, Millikan, Siegbahn, and Compton. By the mid 1930s, however, a mistake was found in Millikan's classic elementary charge measurement due to an error in the value for the viscosity of air.

Once this was corrected the accepted value of the Avogadro constant dropped by 0.6% [5].

References

1. Perrin JB (1965) Discontinuous structure of matter. In: Nobel lectures, Physics. Elsevier, Amsterdam, pp 1922–1941
2. Holden NE (2004) Chem Int 26:4–7
3. Jensen WB (2010) J Chem Educ (in press)
4. Bassow H (1991) J Chem Educ 68: 273-274
5. Lipson H, Riley DP (1943) Nature 151: 250-250