NRC Publications Archive Archives des publications du CNRC

Condensation cascades and methylgroup transfer reactions during the formation of arsane, methyl- and dimethylarsane by aqueous borohydride and (methyl) arsenates

D'Ulivo, Alessandro; Meija, Juris; Mester, Zoltán; Pagliano, Enea; Sturgeon, Ralph E.

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

Publisher's version / Version de l'éditeur:

<https://doi.org/10.1007/s00216-011-5503-4>

Analytical and Bioanalytical Chemistry, 402, 2, pp. 921-933, 2011-10-30

NRC Publications Record / Notice d'Archives des publications de CNRC:

<https://nrc-publications.canada.ca/eng/view/object/?id=906fc520-b4ee-4e69-9ab0-5defc7ccd1fe>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=906fc520-b4ee-4e69-9ab0-5defc7ccd1fe>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

Questions? Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

Vous avez des questions? Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.

Condensation cascades and methylgroup transfer reactions during the formation of arsane, methyl – and dimethylarsane by aqueous borohydride and (methyl) arsenates

Alessandro D’Ulivo^{a*}, Juris Meija^b, Zoltán Mester^b, Enea Pagliano^c, and Ralph E. Sturgeon^b

^a CNR, Consiglio Nazionale delle Ricerche, Istituto di Chimica dei Composti Organometallici, Via G. Moruzzi,1, Pisa, 56124, Italy

^b Institute for National Measurement Standards, National Research Council Canada, Ottawa, ON K1A 0R6, Canada

^c Scuola Normale Superiore, Piazza dei Cavalieri, 7, Pisa, 56126, Italy

*Corresponding author, email: dulivo@pi.iccom.cnr.it

ABSTRACT

The formation of volatile products during the reaction of As(III), As(V), MeAsO(OH)₂ and Me₂AsO(OH) with aqueous NaBH₄ has been investigated and the formation of arsanes, di- and triarsanes has been detected. The presence of triarsanes is reported here for the first time. Di- and triarsanes are likely formed in condensation cascade reactions, whereas trimethylarsane arises *via* the transfer of a methyl group. The formation of volatile byproducts is considerably reduced by increasing the acidity of the medium and the concentration of NaBH₄ or by the addition of thiols, such as cysteine. A reaction scheme is proposed which reconciles the evidence reported herein and elsewhere in the literature that is valid for both analytical (trace analysis) and non-analytical reaction conditions.

Keywords: arsenic, monomethyarsonic acid, dimethylarsinic acid, tetrahydridoborate, diarsanes, triarsanes.

Graphic Abstract

... Condensation cascade

... Methyl transfer

INTRODUCTION

Chemical generation of volatile species (CGVS) of metallic or organometallic compounds by reactions with aqueous NaBR_4 ($R = \text{H}$, alkyl or phenyl group) or XBH_3 ($X = \text{NH}_3$, NR_3 , CN^-) coupled with atomic or molecular spectroscopy or with mass spectrometry is among the most powerful analytical tools for trace element determination and speciation [1-3]. Despite the widespread analytical application of volatile hydride formation for determination of trace amounts of elements such as arsenic, antimony, bismuth, germanium, tin, lead, selenium, tellurium and mercury the mechanism by which they are formed has been debated for many years. In particular, perpetuation of the notorious two-century-old incorrect hypothesis that free atomic (nascent) hydrogen is somehow involved in this reaction needs to be dispelled [4-5].

Recent studies have shown that the aqueous-phase reaction of NaBH_4 with oxoacids species of elements such as arsenic occurs through a concerted (direct) transfer of hydrogen atoms from boron to the element, thus forming the hydride, AsH_3 [6]. Generation of EH_n -type hydrides is a n -step process wherein each of the n hydrogen atoms in the hydride originate from different borane molecules [7]. In addition, acid hydrolysis of NaBH_4 is not necessary for the formation of hydrides [8-9]. Analyte-borane intermediates have also been detected, as in the case of the reaction of $\text{Me}_2\text{AsO}(\text{OH})$ with aqueous aminoboranes [7]. Certain aspects of generation of the volatile hydrides, however, are not fully understood. An additional problem that arises from the incomplete understanding of this reaction is the assumption that the reaction of NaBH_4 with aqueous ions of the hydride-forming elements leads to the formation of a *unique* volatile hydride. It is well-known, however, that with the increased concentration of the analytical substrate, many elements are partially converted to solid reaction products [10-12]. This diminishes the yield of volatile hydride and leads to a curvature in calibration plots at higher concentrations of analyte [11-12]. At trace levels (sub-mg/L), inorganic As(III) and As(V), $\text{MeAsO}(\text{OH})_2$ and $\text{Me}_2\text{AsO}(\text{OH})$ react with an excess of aqueous NaBH_4 , forming AsH_3 , MeAsH_2 and Me_2AsH , respectively [2]. As the concentration of analytical substrate increases, or the amount ratio of NaBH_4 -to-analyte decreases, various volatile byproducts, such as diarsanes or Me_3As , are formed [7]. This phenomenon is of great importance to analytical chemists. For example, it casts doubt on whether diarsane and monomethyl diarsane are indeed present in the environment, as recently reported [13], or they only appear as artefacts of the analytical method, as will be demonstrated here.

This study is devoted to a clarification of the mechanism of arsane generation from inorganic and methyl arsenates under analytical and non-analytical conditions. A series of experiments was performed with deuterium-labeled reagents using gas chromatography coupled with mass

spectrometry for identification of the reaction products. The mass spectra of the hydrides were mathematically deconvoluted in order to obtain the amount fractions of isotopologues $\text{AsH}_n\text{D}_{3-n}$ ($n = 0-3$), $\text{MeAsH}_n\text{D}_{2-n}$ ($n = 0-2$), $\text{Me}_2\text{AsH}_n\text{D}_{1-n}$ ($n = 0-1$), and the amount fraction of deuterium incorporated into the final hydrides [16].

EXPERIMENTAL SECTION

Instrumentation. A Hewlett-Packard 6890 gas chromatograph (GC) operated in splitless mode and equipped with a Hewlett-Packard 5973 mass selective detector (MS) was fitted with a 30 m capillary GC column (Valcobond VB-1; 1 μm film thickness, 0.25 mm internal diameter) for the acquisition of mass spectra of arsane and methylated arsanes isotopologues. The GC was operated under the following conditions: injector temperature: 160 $^{\circ}\text{C}$; oven temperature program: 35 $^{\circ}\text{C}$, hold for 4 min, heated to 200 $^{\circ}\text{C}$ at 15 $^{\circ}\text{C min}^{-1}$. Initial transfer line temperature was 150 $^{\circ}\text{C}$, heated at 30 $^{\circ}\text{C min}^{-1}$ to 250 $^{\circ}\text{C}$. For the separation and identification of the less volatile arsenic species a different GC capillary column was employed (DB-5; 30 m length, 5 μm film thickness, 0.25 mm internal diameter) and it was operated under the following conditions: injector temperature: 160 $^{\circ}\text{C}$; oven temperature program: 35 $^{\circ}\text{C}$, hold for 4 min, heated to 200 $^{\circ}\text{C}$ at 15 $^{\circ}\text{C min}^{-1}$. Initial transfer line temperature was 150 $^{\circ}\text{C}$, heated at 30 $^{\circ}\text{C min}^{-1}$ up to 250 $^{\circ}\text{C}$. Gas-tight syringes (Hamilton; 1 mL and 5 mL) were employed for sampling gases from reaction vial headspace. Screw cap vials (4 mL) fitted with PTFE/silicone septa (borosilicate glass, Pierce Chemical Co.) were used. The carrier gas was He at 1.2 mL min^{-1} . Mass spectral deconvolution of results was undertaken.

Chemicals and procedures. NaBH_4 pellets (Alfa Aesar, Word Hill, MA); NaBD_4 powder (isotopic purity $x_D = 99\%$; Cambridge Isotope Laboratories, MA); 37% DCl in D_2O ($x_D = 99.5\%$; Aldrich); 30% NaOD in D_2O ($x_D = 99\%$; Aldrich) and D_2O ($x_D = 99.9\%$; Aldrich) were used. Arsenic standard solutions (2000 $\mu\text{g As mL}^{-1}$) were prepared from K_3AsO_4 (Baker's Analyzed, >99.0% purity), $\text{MeAsO}(\text{ONa})_2$ (ChemService, PA, USA, 99.0% purity) and cacodylic acid, $\text{Me}_2\text{AsO}(\text{OH})$ (Sigma Aldrich, 99.3% purity). A 1 mol L^{-1} bromide/0.2 mol L^{-1} bromate solution was prepared from analytical grade solid reagents (Carlo Erba). L-Cysteine solution ($\gamma = 10 \text{ g L}^{-1}$) was prepared by dissolving the solid reagent (Fluka) in 0.1 M HCl . A hydrazine solution ($w = 250 \text{ mg g}^{-1}$, $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$; Fluka), was employed without further dilution. All other chemicals were reagent grade. Stock solutions of 1 M NaBH_4 or NaBD_4 , prepared in H_2O , were stabilized with 0.5 M NaOH . A stock solution of 0.2 M NaBD_4 was prepared in D_2O and stabilized with 0.1 M NaOD . All NaBH_4 or NaBD_4 solutions were stored at 4 $^{\circ}\text{C}$ and were stable for a week. Working solutions for reactions were prepared by dilution of the stocks just before use.

Generation of arsanes. For the generation of different volatile arsanes, 1 mL of HCl solution (0.01 – 0.20 M) was placed into a 4 mL reaction vial and spiked with the required amount of the

arsenic standard (5 – 400 μL of 2000 $\mu\text{g As mL}^{-1}$). The vial was sealed, purged with nitrogen and kept under magnetic stirring. An aliquot (0.1–1.0 mL) of 0.2 M NaBH_4 (or NaBD_4) in 0.1 M NaOH (or NaOD) was then added with a syringe. After the addition of the reducing agent, the headspace of the vial (ca. 0.25 mL) was sampled at different reaction times (from 5 s to 20 min) and analyzed by GC–MS.

For H/D exchange experiments, pure hydrogenated or deuterated arsanes were prepared shortly before their use from fully hydrogenated or deuterated media following the above procedure. After reaction, 5 mL of headspace gas was sampled and then injected into another vial containing the exchange solution, H_2O or D_2O , at the required acidity. The vial was continuously shaken and the headspace (ca. 0.25 mL) was periodically sampled over an interval of 5 min.

Arsenic remaining in the liquid phase. Solutions resulting from the reaction of 10 mM As(III) or MeAsO(OH)_2 or $\text{Me}_2\text{AsO(OH)}$ in 1.4 mL 0.1 M HCl with 0.2 mL 0.2 M NaBH_4 in 0.1 M NaOH were analyzed following the same procedure described above for arsanes. During the reaction, several changes are observed for the different arsenic compounds, these are illustrated in Fig. 1. After 30 min, the headspace of the reaction vial was flushed with nitrogen in order to remove volatile products. The solution was analyzed for both the arsenic dissolved in the liquid phase and for the total arsenic (liquid phase + solid phase).

For the determination of arsenic dissolved in the liquid phase, 0.05 mL of the solution was diluted to 100 mL with 0.1 M HCl and 0.1 M cysteine solution. After 30 min, the solution was analyzed using arsane generation with detection by atomic absorption spectrometry (HG–AAS) [14] For the determination of total arsenic, the solution containing the precipitate was acidified with 0.8 mL of 12 M HCl and then bromine was developed *in situ* by the addition of 10 μL of a bromide/bromate solution. The solution was heated at 95 $^\circ\text{C}$ for 1 h in a heating block and, after cooling, the bromine was reduced with 10 μL of hydrazine solution. The determination of arsenic was achieved by HG–AAS, following the same procedure described for dissolved arsenic.

Amount fraction of arsane isotopologues. Separation of the arsane isotopologues $\text{AsH}_n\text{D}_{3-n}$ ($n = 0–3$), $\text{MeAsH}_n\text{D}_{2-n}$ ($n = 0–2$), and $\text{Me}_2\text{AsH}_n\text{D}_{1-n}$ ($n = 0–1$) cannot be accomplished using conventional capillary GC. Their separation, however, can be accurately performed in the mass domain since their mass spectra are different. For dimethylarsane, estimation of the Me_2AsH and Me_2AsD amount fraction can be performed from the fragments at m/z 106 and 107, respectively. Isotopologues of arsane and monomethylarsane, however, cannot be treated in a similar manner since the electron impact mass spectra of these hydrides overlap substantially. Additionally, there

are no experimental strategies for obtaining pure AsH_2D , AsHD_2 or MeAsHD . In this case, relative abundances of the arsane and monomethylarsane isotopologues were estimated using mathematical deconvolution techniques previously developed for quantitation of hydrides EH_2 ($\text{E} = \text{O}, \text{Se}$) [15], EH_3 ($\text{E} = \text{As}, \text{Sb}, \text{Bi}$) and EH_4 ($\text{E} = \text{Ge}, \text{Sn}$) [16] Electron impact mass spectra of the pure arsane isotopologues are given below:

$\text{AsH}_n\text{D}_{n-3}$, $m/z = 75\text{--}81$: 0.145, 0.420, 0.105, 0.328, 0.000, 0.000, 0.000 ($n = 3$); 0.139, 0.262, 0.177, 0.079, 0.343, 0.000, 0.000 ($n = 2$); 0.135, 0.086, 0.324, 0.006, 0.103, 0.346, 0.000 ($n = 1$) and 0.136, 0.000, 0.431, 0.000, 0.088, 0.000, 0.345 ($n = 0$) [16].

$\text{MeAsH}_n\text{D}_{n-2}$, $m/z = 90\text{--}94$: 0.481, 0.079, 0.434, 0.006, 0.000 ($n = 2$); 0.434, 0.151, 0.026, 0.389, 0.000 ($n = 1$) and 0.350, 0.172, 0.068, 0.018, 0.392 ($n = 0$).

$\text{Me}_2\text{AsH}_n\text{D}_{n-1}$, $m/z = 106\text{--}107$: 1.000, 0.000 ($n = 1$) and 0.000, 1.000 ($n = 0$).

Figure 1. Visual change occurring during the reduction of As(III) (a-c) and $\text{Me}_2\text{AsO(OH)}$ (d-f) with NaBH_4 (Color change for As(III) , As(V) and MeAsO(OH)_2 is indistinguishable). Images are taken approx. 1 s, 1 min and 1 h after mixing the reagents, respectively. Experimental conditions: 1.4 mL of As solution (10 mM As in 0.1 M HCl) are reduced by 0.2 mL 0.2 M NaBH_4 in 0.1 M NaOH.

RESULTS AND DISCUSSION

H/D exchange of hydridic hydrogen in methylarsanes. As reported recently, neither AsH_3 nor AsD_3 undergoes H/D exchange when in contact with acidic solutions ($0 < \text{pH}$, $\text{pD} < 7$) for several minutes [17]. In contrast, methylarsanes undergo fast H/D exchange in acidic media [18]. Both MeAsH_2 and MeAsD_2 undergo rapid H/D exchange in strongly acidic or alkaline conditions; however, no significant H/D exchange is observed in the interval of $6 < \text{pH}$, $\text{pD} < 10$ for contact times with the liquid phase of up to 30 s, as evident from Fig. 2. Generation of MeAsD_2 at a final $\text{pH} = 9$ results in a 95% isotopic purity of MeAsD_2 with only 4% of MeAsHD and 1% MeAsH_2 . With the objective of evaluating the mechanism of hydrogen transfer, this was considered a tolerable level of H/D exchange. Both Me_2AsH and Me_2AsD undergo H/D exchange in strongly acidic media (pH , $\text{pD} = 1$), yet are stable at $6 < \text{pH}$, $\text{pD} < 13$ where no significant exchange ($< 8\%$) takes place in the first 5 min. However, pure Me_2AsD could be obtained only in a strongly basic medium ($\text{pH} > 12$), as shown in Fig. 3. In order to avoid any H/D exchange with the solvent, a final $\text{pH} = 13$ was considered more appropriate, even though dimethylarsane is generated quite slowly under these conditions.

The identification of the pH regions where the arsanes are stable against H/D exchange with the media ($\text{HCl}/\text{H}_2\text{O}$) allowed development of experimental designs in which the integrity of the isotopic composition of the generated volatile arsanes could be maintained. This, in turn, allowed probing of the mechanism of the hydride generation.

Mechanism of the hydrogen transfer from borohydride. Arsenic hydride generation experiments with pure NaBD_4 or from mixtures of NaBD_4 and NaBH_4 are useful for deciphering the origin of the hydrogen and, more importantly, the nature of the hydrogen transfer from the borohydride to arsenic. Most notably, it is among the simplest of means to dismiss the atomic hydrogen hypothesis of arsane generation [7]. Experiments with NaBH_4 and NaBD_4 mixtures can also differentiate between the single- and multi-step transfer of hydrogen atoms from borohydride as these two scenarios result in different patterns of arsane isotopologues, as evident in Fig. 4.

In the multi-step hydrogen transfer, the hydrogen atoms in the final hydride arise from different borane molecules whereas in the case of a single-step hydrogen transfer, all hydrogen atoms in the final hydride derive from the same borane molecule. In the reaction of $\text{MeAsO}(\text{OH})_2$ with a 1:1 mixture of $\text{NaBH}_4/\text{NaBD}_4$, the single-step hydrogen transfer would result in two hydrides, MeAsH_2 and MeAsD_2 , both in equal amounts. In contrast, two-step hydrogen transfer would

result in a mixture of MeAsH_2 , MeAsHD and MeAsD_2 in a ratio of 1:2:1, as coefficients in the Pascal triangle. Comparison of the experimental and expected distributions of monomethylarsane isotopologues, shown in Figures 4c and 4d, clearly favors the two-step transfer of hydrogen from boron to arsenic, analogous to the three-step transfer observed in the formation of AsH_3 , SbH_3 or BiH_3 [7].

Figure 2. H/D exchange of monomethylarsane. Pre-formed Me_2AsD (Me_2AsH) was placed in contact with the exchange media under various conditions. It is reported the amount fractions of $\text{MeAsH}_n\text{D}_{(2-n)}$ ($n = 0, 1, 2$) detected in the headspace at different acidity.

- (a) MeAsH_2 in D_2O at different pD, 30 s contact time.
- (b) Same as (a) but with 5 min contact time.
- (c) MeAsD_2 in H_2O at different pH, 30 s contact time.
- (d) Same as (c) but with 5 min contact time.

Isotope effects. The expected isotopologue distributions were calculated without considering the H/D isotope effect. Indeed, there is no evidence of a significant isotope effect in the formation of dimethylarsane (Fig. 4a). In contrast to this, the isotopologue distributions of arsane, (Fig. 4e) and monomethylarsane (Fig. 4c) are slightly skewed towards more favorable incorporation of H over

D, even if it appears of little relevance. Considering that no isotope effect is observed during the formation of dimethylarsane, this skew is likely due to a secondary isotope effect. This secondary isotope effect is reasonably due to fact that the incorporation rate of an incoming H or D is dependent on the coordination sphere of arsenic, let say to the presence of As-H or As-D bonds which are already formed in the preceding transfer step. In sum, the kinetic isotope effect, which has been observed by replacing H with D in NaBH_4 , indicates that hydrogen transfer is not the rate-determining step in the formation of volatile arsanes.

Figure 3. Synthesis of dimethylarsane. 0.1 mM $\text{Me}_2\text{AsO}(\text{OH})$ placed in a completely hydrogenated media was reduced with 0.2 M NaBD_4 . The isotopic composition of dimethylarsane evolved has been followed against the acidity of the resulting solution. The amount fraction of Me_2AsD is reported at different pH.

Figure 4. Experimental (a, c, e) and predicted (b, d, f) distributions of arsane and methylated arsane isotopologues arising from the reduction of the corresponding pentavalent arsenic compounds.

(a) Dimethylarsane. Conditions: 1 mL 60 μM $\text{Me}_2\text{AsO}(\text{OH})$ in 1 mM HCl was reduced with 0.5 mL of 0.2 M ($\text{NaBH}_4 + \text{NaBD}_4$) in 0.1 M NaOH; all reagents were prepared in H_2O . (b) Predicted distribution of dimethylarsane isotopologues based on the direct hydrogen transfer from borohydride to arsenic.

(c) Monomethylarsane. Conditions: 1 mL of 60 μM $\text{MeAsO}(\text{OH})_2$ in 0.1 M HCl was reduced with 0.5 mL of 0.2 M ($\text{NaBH}_4 + \text{NaBD}_4$) in 0.1 M NaOH; all reagents were prepared in H_2O . (d) Predicted distribution of monomethylarsane isotopologues based on the direct hydrogen transfer from borohydride to As: two-step hydrogen transfer.

(e) Arsane. Conditions: 1 mL of 60 μM $\text{AsO}(\text{OH})_3$ in 1 M HCl was reduced with 0.5 mL of 0.2 M ($\text{NaBH}_4 + \text{NaBD}_4$) in 0.1 M NaOH; all reagents were prepared in H_2O . (f) Predicted distribution of arsane isotopologues based on the direct hydrogen transfer from borohydride to As: three-step hydrogen transfer.

Volatile reaction byproducts. Under analytical conditions and at ultra-trace levels of arsenic, hydride generation is commonly assumed to yield a single volatile product: AsH_3 from As(III) and As(V) , MeAsH_2 from MeAsO(OH)_2 and Me_2AsH from $\text{Me}_2\text{AsO(OH)}$. To test this assumption, various experiments were conducted under analytical and non-analytical conditions wherein the concentration of the arsenic substrates was varied in the range from 10^{-2} to 10 mM (the typical analytical working range is from 1 nM to 1 μM of arsenic). While at the ultra-trace level, the arsenic compounds are indeed converted to unique volatile arsanes, as mentioned above, the formation of other volatile arsanes is evident at higher analyte concentration, as summarized in Tables 1 and 2. In certain cases, these byproducts even become the major components of the headspace.

At arsenic concentrations up to 1 mM, the formation of a reddish suspended particulate matter is also evident for inorganic As and MeAsO(OH)_2 , suggesting that they are not completely converted into volatile species (Fig. 1c). For $\text{Me}_2\text{AsO(OH)}$, no particulate matter is formed, but a white gas is generated upon addition of NaBH_4 and it is clearly visible in the headspace of the reaction vial (Fig. 1e). At the highest concentration level of arsenic (10 mM), quantitative experiments indicate that a large fraction of arsenic remains in the vial after reaction with NaBH_4 . For the inorganic arsenic compounds, about 65% of the total As remains in the condensed phase (45% in solution and 20% as solid precipitate). For both MeAsO(OH)_2 and $\text{Me}_2\text{AsO(OH)}$, the amount fraction of the total As remaining in solution as non-volatile compounds is greater than 98%, but in the latter no precipitate is visible.

In spite of the small fraction amount of volatile species that are formed under these reaction conditions, their identification can give interesting information about the reactions which took place in the liquid phase.

For inorganic arsenic, the only observed volatile byproducts are diarsane ($\text{H}_2\text{As-AsH}_2$), and triarsane ($\text{H}_2\text{As-AsH-AsH}_2$). A gamut of volatile compounds was produced during reduction of methylated arsenic acids by NaBH_4 , including dimeric and trimeric arsenic species. A summary of dimeric and trimeric arsenic species which have been detected in the headspace is reported in Table 4. Mass spectra of trimeric species are reported in Figures 5–9.

A rather interesting behavior is exhibited by the reaction of $\text{Me}_2\text{AsO(OH)}$ with NaBH_4/HCl , which under certain conditions can give Me_3As as the most abundant volatile product (Table 2). Experimental evidence indicates that Me_3As is formed simultaneously with the disappearance of Me_2AsH . In general, formation of volatile byproducts is more pronounced with increased reaction time, As/NaBH_4 amount ratio or acidity. It is interesting to note that the reaction of $\text{Me}_2\text{AsO(OH)}$ with metallic zinc in HCl generates the same volatile reaction products as with NaBH_4 (Table 3).

The addition of cysteine (R–SH) suppressed the formation of unwanted byproducts (Tables 1–2). In particular, it was rather effective for $\text{Me}_2\text{AsO(OH)}$ where the formation of Me_3As was completely eliminated (Table 2).

It is well-known that thiols reduce arsenic compounds to thiolates such as $\text{As}(\text{SR})_3$, $\text{MeAs}(\text{SR})_2$ and $\text{Me}_2\text{As}(\text{SR})$ [19,20,21,22], which are quantitatively converted by NaBH_4 into the corresponding hydrides [23,24]. Consequently, cold vapor generation by NaBH_4 in the presence of thiols presents an attractive way to estimate the purity of the methylarseno compounds.

Figure 5. EI mass spectrum of As_3H_5 . Molecular ion at $m/z = 230$ Da $[\text{As}_3\text{H}_5]^+$. Fragment ions: m/z 225 $[\text{As}_3]^+$; m/z 152 $[\text{As}_2\text{H}_2]^+$; m/z 151 $[\text{As}_2\text{H}]^+$; m/z 150 $[\text{As}_2]^+$; m/z 75 $[\text{As}]^+$.

Figure 6. EI mass spectrum of MeAs_3H_4 . Molecular ion at $m/z = 244$ Da $[\text{MeAs}_3\text{H}_4]^+$. Fragment ions: m/z 225 $[\text{As}_3]^+$; m/z 166 $[\text{MeAs}_2\text{H}]^+$; m/z 150 $[\text{As}_2]^+$; m/z 89 $[\text{CH}_2\text{As}]^+$; m/z 75 $[\text{As}]^+$.

Figure 7. EI mass spectrum of $\text{Me}_2\text{As}_3\text{H}_3$. Molecular ion at $m/z = 258$ Da $[\text{Me}_2\text{As}_3\text{H}_3]^+$. Fragment ions: m/z 241 $[\text{MeAs}_3\text{H}]^+$; m/z 225 $[\text{As}_3]^+$; m/z 181 $[\text{Me}_2\text{As}_2\text{H}]^+$; m/z 165 $[\text{MeAs}_2]^+$; m/z 151 $[\text{As}_2\text{H}]^+$; m/z 105 $[\text{Me}_2\text{As}]^+$; m/z 90 $[\text{MeAs}]^+$; m/z 75 $[\text{As}]^+$.

Figure 8. EI mass spectrum of $\text{Me}_3\text{As}_3\text{H}_2$. Molecular ion at $m/z = 272$ Da $[\text{Me}_3\text{As}_3\text{H}_2]^+$. Fragment ions: m/z 255 $[\text{Me}_2\text{As}_3]^+$; m/z 225 $[\text{As}_3]^+$; m/z 180 $[\text{Me}_2\text{As}_2]^+$; m/z 165 $[\text{MeAs}_2]^+$; m/z 150 $[\text{As}_2]^+$; m/z 105 $[\text{Me}_2\text{As}]^+$; m/z 89 $[\text{CH}_2\text{As}]^+$; m/z 75 $[\text{As}]^+$.

Figure 9. EI mass spectrum of Me_3As_3 . Molecular ion at $m/z = 270$ Da $[\text{Me}_3\text{As}_3]^+$. Fragment ions: m/z 255 $[\text{Me}_2\text{As}_3]^+$; m/z 225 $[\text{As}_3]^+$; m/z 180 $[\text{Me}_2\text{As}_2]^+$; m/z 165 $[\text{MeAs}_2]^+$; m/z 150 $[\text{As}_2]^+$; m/z 105 $[\text{Me}_2\text{As}]^+$; m/z 89 $[\text{CH}_2\text{As}]^+$; m/z 75 $[\text{As}]^+$.

Figure 10. EI mass spectrum of $\text{H}_2\text{As-SbH}_2$. Molecular ion at $m/z = 202$ Da $[\text{H}_2^{75}\text{As-}^{123}\text{SbH}_2]^+$ and 200 Da $[\text{H}_2^{75}\text{As-}^{121}\text{SbH}_2]^+$. Fragment ions: m/z 198 $[\text{H}_2^{75}\text{As-}^{123}\text{Sb}]^+$ and m/z 196 $[\text{H}_2^{75}\text{As-}^{121}\text{Sb}]^+$.

Table 1. Volatile arsenic compounds detected during the reaction of inorganic As^{III}, As^V or MeAsO(OH)₂ with NaBH₄

Compound	As /NaBH ₄ μmol/μmol	Conditions ^a	Time (min)	Volatile arsenic species ^b							
				AsH ₃	MeAsH ₂	Me ₂ AsH	Me ₃ As	Me ₂ AsCl	As ₂ H ₄	H ₂ As ₂ MeH	As ₂ (HMe) ₂
AsO(OH) ₃	2.7/20	pH = 3	0.2	100	–	–	–	–	9.5	–	–
	2.7/100	pH = 9	0.2	100	–	–	–	–	–	–	–
As(OH) ₃	2.7/20	pH = 3	0.2	100	–	–	–	–	1.4	–	–
	2.7/100	pH = 9	0.2	100	–	–	–	–	0.2	–	–
MeAsO(OH) ₂	10.8/8	pH = 3–4	0.2	–	100	–	–	–	–	–	–
			30	–	–	–	–	–	100	–	–
	2.7/20	pH = 3	0.2	–	–	–	–	2.1	100	0.2	4.9
			20	7.4	100	0.6	0.3	–	–	–	–
	1.3/100	pH = 9	0.2	1.3	100	0.14	–	–	–	–	0.83
			0.2	2.4	100	0.18	–	–	–	–	3.5
	2.7/100	pH = 9	0.2	0.16	100	0.14	–	–	–	–	0.6
		0.2 M Cys	20	0.14	100	0.37	–	–	–	–	0.27
		pH = 6	0.2	0.17	100	0.14	–	–	–	–	0.35
		0.2 M Cys	20	0.21	100	0.35	–	–	–	–	–
	0.27/100	pH = 9	0.2	0.1	100	–	–	–	–	–	–
		0.2 M Cys	0.2	0.1	100	–	–	–	–	–	–

a – 0.5 mL NaBH₄ added to 1 mL sample solution. Final pH is given.

b – Relative signals are the integrated area of the TIC peaks. Relative signals < 0.1 are not reported.

Table 2. Volatile arsenic compounds detected during the reaction $\text{Me}_2\text{AsO}(\text{OH})$ with NaBH_4

As / NaBH_4 $\mu\text{mol}/\mu\text{mol}$	Conditions ^a	Time (min)	Volatile arsenic species ^b								
			AsH_3	MeAsH_2	Me_2AsH	Me_3As	Me_2AsO	Me_2AsOH	Me_2AsCl	$\text{As}_2(\text{O})\text{Me}_4$	As_2Me_4
0.27/100	pH = 3	0.2	–	8.1	100	7.1	–	–	–	–	–
		20	–	15.1	6.9	100	–	–	–	–	–
0.13/100	pH = 9	5	0.3	4.32	100	9.0	–	–	–	–	–
	pH = 13	5	–	–	100	–	–	–	–	–	–
2.7/20	pH = 3	4	0.1	0.1	1.5	100	–	0.2	2.1	0.13	0.2
		20	–	–	–	100	–	–	0.5	–	0.3
0.27/100	pH = 9	0.2	–	4	100	2.9	–	–	–	–	0.9
		20	26	11	59	100	–	–	–	–	–
10.8/8	pH = 2	0.2	–	1.3	–	100	0.7	1.1	0.9	5.6	5.0
		30	–	–	–	100	0.2	7.9	1.5	9.4	4.2
	pH = 4	20	–	–	–	100	0.2	6.4	1.3	8.5	5.4
		20	–	–	16	100	0.6	2.2	0.9	9.8	7.0
0.27/100	pH = 3 0.2 M Cys	0.2	4	0.3	100	–	–	–	–	–	–
		20	–	0.3	100	0.1	–	–	–	–	–
	pH = 9 0.2 M Cys	0.2	–	0.1	100	–	–	–	–	–	–
		20	–	0.1	100	0.1	–	–	–	–	–
0.4/100	pH = 9 0.2 M Cys	0.2	–	0.2	100	–	–	–	–	–	–

a – 0.05 – 0.5 mL NaBH_4 added to 1 mL sample solution. Final pH is given.

b – Relative signals are the integrated area of the TIC peaks. Relative signals < 0.1 are not reported.

Table 3. Volatile arsenic compounds detected during $\text{Me}_2\text{AsO}(\text{OH})$ reaction with metallic Zn powder

Conditions ^a	Time (min)	Volatile arsenic species ^b							
		AsH_3	MeAsH_2	Me_2AsH	Me_3As	Me_2AsOH	Me_2AsCl	As_2Me_4	
1 M HCl	2	–	3.7	100	13.4	2.4	0.6	7.3	
	25	3.9	3.5	100	38.5	0.1	–	0.4	
0.1 M HCl	2	–	1.0	100	0.64	0.1	–	0.6	
	25	12.5	5.4	76	100	–	–	0.3	
	65	11.4	5.2	51	100	–	–	–	

a – 1.1 μmol $\text{Me}_2\text{AsO}(\text{OH})$ into 1 mL sample solution reacted by 150 μmol Zinc.

b – Relative signals are the integrated area of the TIC peaks. Relative signals < 0.1 are not reported.

Table 4. Volatile di- and triarsanes detected in the headspace of reaction vials

As substrate	Arsanes (a)	Diarsanes		Triarsanes		Mass Spectra
		t_R / min (a)	t_R / min (b)	t_R / min (b)		
As(III)	AsH ₃	1.35	H ₂ As–AsH ₂	2.64	H ₂ As–AsH–AsH ₂	Fig. 7
	MeAsH ₂	1.43	MeHAS–AsH ₂	4.45	Me ₃ As ₃ H ₂	Fig. 10
AsO(OH) ₂			MeHAS–AsHMe	6.60	Me ₃ As ₃	Fig. 11
			Me ₂ As–AsHMe	7.30		
	Me ₂ AsH	1.65	Me ₂ As–AsHMe	7.30		
Me ₂ AsO(OH)	Me ₃ As	1.84	Me ₄ As ₂ O	7.81		
			Me ₄ As–AsMe ₂	8.06		
MeAsO(OH) ₂ and As(III)	AsH ₃	1.35	H ₂ As–AsH ₂	2.64	MeAs ₃ H ₄	Fig. 8
	Me ₂ AsH	1.65	MeHAS–AsH ₂	4.45	Me ₂ As ₃ H ₃	Fig. 9
			Me ₂ As–AsH	5.35		
			MeHAS–AsHMe	6.60		
MeAsO(OH) ₂ and Me ₂ AsO(OH)	AsH ₃	1.35	Me ₂ As–AsH ₂	5.35	Me ₃ As ₃	Fig. 11
	Me ₂ AsH	1.65	Me ₂ As–AsHMe	6.60		
	Me ₃ As	1.84	Me ₂ As–AsMe ₂	8.06		

In all proofs an acid solution (0.1 M HCl) containing the above mentioned As compounds (10 mM As content) was reduced by 0.2 M NaBH₄ in 0.1 M NaOH. A final pH = 3–4 was measured.

(a) Retention times of some detected monoarsanes is given for comparison of retention times with di- and triarsanes. Other detected monoarsanes are reported in Table 2.

(b) Retention times obtained with DB-5 column. See Instrumentation section for details.

Mechanism of volatile byproduct formation: condensation reactions. The formation of volatile byproducts is a result of the interaction between the reduction intermediates and the volatile arsanes. For example, pure Me₂AsH (prepared separately from MeAsO(OH)₂/NaBH₄ in the presence of cysteine), when injected into a vial containing aqueous Me₂AsO(OH), does not result in formation of Me₃As nor diarsanes. According to the evidence collected herein, it can be assumed that the stepwise formation of arsanes proceeds through a series of intermediates, as summarized in schemes 2–5:

Here, the reagents and all identified compounds are shown in bold. In each of the reaction sequences (1–4), arsenic becomes progressively less reactive towards transfer of hydrogen from boron as a result of the decreasing number of oxygen atoms bound to it. The intermediates involved in the last hydrogen transfer step are, therefore, the least reactive towards the hydride attack in their relevant reaction sequences. Likewise, the reactivity towards transfer of hydrogen from NaBH₄ is expected to decrease in the order AsH₂(OH) < MeAsH(OH) < Me₂AsOH. This is due to the replacement of the –OH groups by the methyl groups. The simultaneous presence of the reaction intermediates and the final hydrides leads to the formation of diarsanes *via* condensation reactions:

The bidirectional reaction (5) is also supported by Rheingold et al. [25] who reported that tetramethyl diarsane was formed in a reaction of Me₂AsH with Me₂AsX (X = Cl, Br, I or CN), in benzene:

When aqueous As(III), MeAsO(OH)₂ or Me₂AsO(OH), alone or in combination with each other, are treated with NaBH₄, all (six) possible structural isomers of diarsanes were detected in the headspace, *i.e.*, H₂As–AsH₂, MeHAs–AsH₂, Me₂As–AsH₂, MeHAs–AsHMe, Me₂As–AsHMe and Me₂As–AsMe₂, including the oxidation product (Me₂As–As(O)Me₂) of the latter and some triarsane species (Table 4). This clearly indicates that a condensation cascade took place among all possible intermediates which are formed in the reaction pathways of different arsenic substrates, including the final hydrides. The interaction between the proposed reaction intermediates is also confirmed in experiments where As(III) was reduced in the presence of Sb(III). In this case the formation of the dimer H₂As–SbH₂ was observed (Fig. 10). The stibodiarsane could be formed only by reaction of AsH₃ with SbH₂OH or AsH₂OH with SbH₃.

The formation of di- and triarsanes also provides an explanation for the formation of precipitates in the case of As(III), As(V) and MeAsO(OH)₂. When mono-arsenic species contain either only one –H or –OH moiety, the condensation cascade terminates with the formation of a dimer (see

reactions 5-7). Triarsanes, on the other hand, can be formed by condensation reactions of the following type:

In cases when reaction intermediates contain arsenic bound to hydride and hydroxo groups ($\text{H}-\text{AsX}-\text{OH}$, $\text{X} = \text{Me}, \text{H}$ or OH), the condensation reaction may proceed to the formation of linear or branched polymers,

or cyclic As species,

that remain in the solid phase.

The reaction schemes presented above are in agreement with observations reported elsewhere. Under conditions typically employed in analytical protocols for trace analysis, the main reaction products are the simple volatile arsanes (reactions 1-4). Approaching conditions for synthesis reactions, however, the occurrence of condensation reactions become more pronounced, which lead to the formation of non-volatile arsenic species that remain in the reaction solution in either soluble or insoluble form. For example, reddish solid arsenic hydrides, arising from chemical reduction of inorganic As(III) in aqueous solution, are reported by Jolly et al. [26]. In the case of $\text{MeAsO}(\text{OH})_2$, its reduction with H_3PO_2 resulted in the formation of linear or cyclic methylpolyarsanes [27,28]. Under similar conditions, reduction of $\text{Me}_2\text{AsO}(\text{OH})$ with H_3PO_2 yields the water-soluble $\text{Me}_2\text{As}-\text{AsMe}_2$ [27]. The fully methylated diarsane cannot react further with other arsenic intermediates from reaction 1 and therefore is the terminus of the condensation cascade.

Formation of trimethylarsane. The observation that trimethylarsane, and not dimethylarsane, is the main volatile product of reduction of $\text{Me}_2\text{AsO}(\text{OH})$ under non-analytical reaction conditions, is puzzling. It appears, from results reported in Table 2, that the formation of Me_3As coincides with the disappearance of Me_2AsH . This indicates that Me_3As is formed by interaction of Me_2AsH with an arsenic intermediate capable of acting as a methylation agent. The presence of such species was confirmed in the reaction of $\text{Me}_2\text{AsO}(\text{OH})$ and NaBH_4 in the presence of Sb(III) . The formation of MeSbH_2 was confirmed. The formation of Me_3As during reduction of $\text{Me}_2\text{AsO}(\text{OH})$

by NaBH_4 is likely due to the known intermediate $\text{Me}_2\text{As}-\text{AsMe}_2$. Tetramethyl diarsane is also able to form Me_2AsH from MeAsH_2 [29]. As described above, formation of diarsanes is suppressed when the NaBH_4 reduction is performed in the presence of thiols. When this is conducted with the $\text{Me}_2\text{AsO}(\text{OH})/\text{NaBH}_4$ reaction, Me_3As is not formed (Table 2).

Methyl transfer reactions take place also in the case of elevated concentration of $\text{MeAsO}(\text{OH})_2$ as it result evident from the results reported in Table 1 (formation of AsH_3 , Me_2AsH) and in Table 4 (formation of $\text{MeHAs}-\text{AsH}_2$ and $\text{MeHAs}-\text{AsMe}_2$ cannot be explained only by condensation reactions). The effects produced by methyl transfer reactions during the reaction of $\text{MeAsO}(\text{OH})_2$ with NaBH_4/HCl are much less pronounced than those observed for $\text{Me}_2\text{AsO}(\text{OH})$, and the main volatile product is still the expected MeAsH_2 (>92%).

CONCLUSIONS

The mechanism for the reaction of arsenates – $\text{As}(\text{OH})_3$, $\text{AsO}(\text{OH})_3$, $\text{MeAsO}(\text{OH})_2$, $\text{Me}_2\text{AsO}(\text{OH})$ – with aqueous tetrahydroborate(III) has been proposed. It applies to a wide range of reaction conditions employed both in trace analysis and synthesis. Under analytical conditions, the action of NaBH_4 on the arsenic substrate promotes their conversion to the corresponding volatile hydrides AsH_3 , MeAsH_2 , and Me_2AsH . This process is the consequence of the direct step-wise hydrogen transfer from tetrahydroborate(III) to arsenic. The formation of volatile hydrides passes through arseno- intermediates wherein the $-\text{OH}$ groups are sequentially replaced by hydrogen atoms. Under non-analytical (synthesis) conditions, reduction by NaBH_4 competes with two other processes: condensation between $\text{As}-\text{H}$ and $\text{HO}-\text{As}$ and transfer of methyl group. The first effect arises from the condensation of intermediates and causes the formation of $\text{As}-\text{As}$ bonds. Consequently, volatile di- and triarsanes are formed along with a reddish precipitate of polymeric arsanes. The transfer of a methyl group between the reaction products and intermediates is peculiar to the reduction of $\text{Me}_2\text{AsO}(\text{OH})$ with NaBH_4 . Here, the identification of Me_3As as the main volatile reduction product is evidence that a transfer of methyl group has occurred. The likely alkylating agent seems to be the condensation cascade product, As_2Me_4 , which has been detected in the headspace of the reaction medium.

The detected cross reactions among intermediates belonging to different reaction pathways of different arsenic compounds and arsenic and antimony compounds, makes the proposed reaction model also interesting for the interpretation and the comprehension of interference mechanisms in CGVS.

REFERENCES

1. Sturgeon RE, Mester Z (2002) Appl Spectrosc 56: 202A-213A
2. Dědina J, Tsalev DL, Hydride Generation Atomic Absorption Spectroscopy, Wiley, Chichester, 1995
3. D'Ulivo A, Loreti V, Onor M, Pitzalis E, Zamboni R (2003) Anal Chem 75: 2591-2600
4. D'Ulivo A (2010) Spectrochim Acta Part B 65: 360-375
5. D'Ulivo A, Dědina J, Mester Z, Sturgeon RE, Wang Q, Welz B (2011) Pure Appl Chem 83:1283-1340
6. D'Ulivo A, Mester Z, Sturgeon RE (2005) Spectrochim Acta Part B 60: 423-438
7. D'Ulivo A, Mester Z, Meija J, Sturgeon RE (2007) Anal Chem 79: 3008-3015
8. D'Ulivo A (2004) Spectrochim Acta Part B 59: 793-825
9. D'Ulivo A, Baiocchi C, Pitzalis E, Onor M, Zamboni R (2004) Spectrochim Acta Part B 59: 471-486
10. Petrick K, Krivan V (1987) Fresenius Z Anal Chem 327: 338-342
11. D'Ulivo A, Marcucci K, Bramanti E, Lampugnani L, Zamboni R (2000) Spectrochim Acta, Part B 55: 1325-1336
12. D'Ulivo A, Battistini SST, Pitzalis E, Zamboni R, Mester Z, Sturgeon RE (2007) Anal Bioanal Chem 388: 783-791
13. Köster J, Diaz-Bone RA, Planer-Friederich B, Rothweiler B, Hirner H (2003) J Mol Structure 661-662: 347-346
14. Pitzalis E, Ajala D, Onor M, Zamboni R, D'Ulivo A (2007) Anal Chem 79: 6324-6333
15. Meija J, Mester Z, D'Ulivo A (2006) J Am Soc Mass Spectrom 17: 1028-1036
16. Meija J, Mester Z, D'Ulivo A (2007) J Am Soc Mass Spectrom 18: 337-345
17. D'Ulivo A, Mester Z, Meija J, Sturgeon RE (2006) Spectrochim Acta Part B 61: 778-787
18. Pergantis SA, Winnik W, Eimar W.R, Cullen WR (1997) Talanta 44: 1941-1947
19. Raab A, Meherag AA, Jaspars M, Genney DR, Feldmann J (2004) J Anal At Spectrom 19: 183-190
20. Rey NA, Howarth OW, Pereira-Maia ECJ (2004) Inorg Biochem 98: 1151-1159
21. Scott N, Hatlelid KM, MacKenzie NE, Carter DE (1993) Chem Res Toxicol 6: 102-106
22. Spuches AM, Kruszynska HG, Rich AM, Wilcox DE (2005) Inorg Chem 44: 2964-2972
23. Le X-C, Cullen WR, Reimer KJ, (1994) Anal Chim Acta 285: 277-285
24. Tsalev DL, Sperling M, Welz B (2000) Talanta, 51, 1059-1068
25. Rheingold AL; Pleau EJ, Ferrar WT (1977) Inorg Chim Acta 22: 215-218
26. Jolly WL, Anderson LB, Beltrami RT. (1957) J Am Chem Soc 79: 2443-2447
27. Knoll F, Marsmann HC, Van Wazer JR (1969) J Am Chem Soc 91: 4986-4989
28. Waser J, Schoemaker V (1945) J Am Chem Soc 67: 2014-2018
29. Gupta VK, Krannich LK, Watkins L (1987) Inorg Chim Acta 132: 163-164