

NRC Publications Archive Archives des publications du CNRC

Multipoles from Majorana constellations

Romero, J. L.; Klimov, A. B.; Goldberg, A. Z.; Leuchs, G.; Sánchez-Soto, L. L.

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. / La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien DOI ci-dessous.

Publisher's version / Version de l'éditeur:

<https://doi.org/10.1103/PhysRevA.109.012214>

Physical Review A, 109, 1, 2024-01-24

NRC Publications Archive Record / Notice des Archives des publications du CNRC :

<https://nrc-publications.canada.ca/eng/view/object/?id=2a232ec7-3d56-4b4b-b801-c26b0ce95728>

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=2a232ec7-3d56-4b4b-b801-c26b0ce95728>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

Questions? Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the first page of the publication for their contact information.

Vous avez des questions? Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.

Multipoles from Majorana constellations

J. L. Romero ¹, A. B. Klimov ², A. Z. Goldberg ^{3,4}, G. Leuchs ^{5,6} and L. L. Sánchez-Soto ^{1,5}

¹*Departamento de Óptica, Facultad de Física, Universidad Complutense, 28040 Madrid, Spain*

²*Departamento de Física, Universidad de Guadalajara, 44420 Guadalajara, Jalisco, Mexico*

³*National Research Council of Canada, Ottawa, Ontario, Canada K1N 5A2*

⁴*Department of Physics, University of Ottawa, Ottawa, Ontario, Canada K1N 6N5*

⁵*Max-Planck-Institut für die Physik des Lichts, 91058 Erlangen, Germany*

⁶*Institut für Optik, Information und Photonik, Friedrich-Alexander-Universität Erlangen-Nürnberg, 91058 Erlangen, Germany*



(Received 22 May 2023; revised 22 September 2023; accepted 2 January 2024; published 22 January 2024)

Majorana stars, the $2S$ spin coherent states that are orthogonal to a spin- S state, offer an elegant method to visualize quantum states, disclosing their intrinsic symmetries. These states are naturally described by the corresponding multipoles. These quantities can be experimentally determined and allow for an $SU(2)$ -invariant analysis. We investigate the relationship between Majorana constellations and state multipoles, thus providing insights into the underlying symmetries of the system. We illustrate our approach with some relevant and informative examples.

DOI: [10.1103/PhysRevA.109.012214](https://doi.org/10.1103/PhysRevA.109.012214)

I. INTRODUCTION

Completely symmetric states aptly describe numerous phenomena [1]. These states play a crucial role in the characterization of spinor Bose gases [2,3], whose dynamics have been observed with spin-1 [4,5], spin-2 [6], and even spin-3 [7,8] condensates. They have also been used for the characterization of entanglement [9–14] in those boson systems.

A salient feature of this family is that any state of spin S can be described as a permutation-symmetric n -qubit state, with $n = 2S$. This is the basis of the elegant representation devised by Majorana [15], in which a spin S is depicted by $2S$ points (called the stars) on the Bloch sphere. Although the subject attracted some attention in relation to the quantum theory of angular momentum [16–19], it remained quiescent until 1969, when Penrose reinterpreted Majorana stars as principal null directions in spinor theory [20] and brought it to wider attention in his celebrated book [21].

Apart from indisputable mathematical advantages [22], this picture builds a bridge between the abstract Hilbert space (where the states live) and the simple geometry of the Bloch sphere. Consequently, this representation rapidly meets the increasing interest in high-dimensional quantum systems and, several decades after its conception, is being used in fields as diverse as polarization [23–27], spinor Bose gases [28–31], multiqubit systems [32–38], metrology [39–43], geometric phases [44–51], non-Hermitian lattices [52], and algebraic quantum models such as the Lipkin-Meshkov-Glick model [53,54].

The distribution of Majorana stars conveys complete information and can be directly computed when the quantum state is known. But this notion has far-reaching advantages: by visualizing in a crystal-clear manner the intrinsic symmetries of the state, one can perceive connections with other intriguing questions. Examples include geometrical measures of entanglement [55–57], spherical t -designs [58,59], and the Thomson [60–65] and Tammes [66–68] problems. Moreover, a number of states with remarkable properties, such as queens [69] and kings of quantumness [70], maximally entangled states [71], k -uniform states [72–74], and states with maximal Wehrl entropy [75], can be aptly understood in terms of the properties of their corresponding constellations [76].

Distributing points on a sphere is a mathematical problem with a long history and with a variety of optimal configurations depending on the cost function one tries to optimize [77–80]. This suggests exploring those arrangements of points distinguished by some extremal properties. Then a natural question arises: if one knows the locations of the Majorana stars, what can one say about the state, in particular, about its multipolar distribution? That is precisely our main goal here.

For a system of point charges on the sphere, the most suitable way of capturing the progressively finer angular features of the system is the standard multipolar expansion [81]. Such an expansion can often be truncated, meaning that to a good approximation only the first terms need to be retained. We propose here to carry out a similar procedure for the Majorana constellation: the resulting multipoles constitute a basic tool for problems with an $SU(2)$ invariance [82,83] and, in addition, they can be experimentally determined with simple procedures [84,85].

The paper is organized as follows. In Sec. II we introduce the basic notions needed to understand the Majorana constellations, whereas in Sec. III we show how to calculate the multipoles from a given Majorana constellations. Our

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Open access publication funded by the Max Planck Society.

insight is to apply the time-honored Vieta formulas [86] to the polynomial defining the constellation, which provides a shortcut between the stars and other representations of the state. In Sec. IV we thoroughly examine how the method works in a series of relevant examples. Further physical implications are discussed in Sec. V and, finally, our conclusions are summarized in Sec. VI.

II. MAJORANA CONSTELLATIONS

We will deal with any pure system living in a finite-dimensional Hilbert space of dimension $2S + 1$, which can be formally regarded as a spin S . The corresponding space \mathcal{H}_S , spanned by the standard angular momentum basis $\{|S, m\rangle \mid m = -S, \dots, S\}$, is the carrier of the irreducible representation (irrep) of spin S of $SU(2)$ and is isomorphic to \mathbb{C}^{2S+1} . Since any two vectors in \mathcal{H}_S differing by a phase represent the same physical state, the manifold of physical states is the projective space $\mathbb{C}P^{2S}$ [87].

The merit of Majorana was to show that points in $\mathbb{C}P^{2S}$ are in one-to-one correspondence with unordered sets of (possibly coincident) $2S$ points on the unit sphere \mathcal{S}_2 . There are various ways to see why this is so, but probably the most direct one is in terms of coherent states.

The spin (or Bloch) coherent states live in \mathcal{H}_S and are displaced versions of a fiducial state, much the same as for the canonical coherent states on the plane. This fiducial state is chosen so as to minimize the variance of the Casimir operator $\mathbf{S}^2 = S_x^2 + S_y^2 + S_z^2$, where (S_x, S_y, S_z) are the angular momentum operators, which generate the algebra $\mathfrak{su}(2)$. The minimum-variance states are $|S, \pm S\rangle$ and they guarantee that their displaced versions are the closest to classical states. The displacement operator on \mathcal{S}_2 is $D(\theta, \phi) = \exp(i\phi S_z) \exp(i\theta S_y) = \exp[\frac{1}{2}\theta(S_+ e^{-i\phi} - S_- e^{i\phi})]$, where $S_{\pm} = S_x \pm iS_y$ are raising and lowering operators. Disentangling this displacement allows us to express the coherent states $|\theta, \phi\rangle = D(\theta, \phi)|S, -S\rangle$ as [88,89]

$$|\theta, \phi\rangle \equiv |z\rangle = \frac{1}{(1 + |z|^2)^S} \exp(zS_+) |S, -S\rangle, \quad (2.1)$$

where the label $z = \tan(\theta/2)e^{-i\phi}$ corresponds to an inverse stereographic projection from the south pole, mapping the point $z \in \mathbb{C}$ onto the point $(\theta, \phi) \in \mathcal{S}_2$ [90].

On expanding the exponential, we can write the coherent states in terms of the basis states of the irrep:

$$|z\rangle = \frac{1}{(1 + |z|^2)^S} \sum_{m=-S}^S \binom{2S}{S+m}^{\frac{1}{2}} z^{S+m} |S, m\rangle, \quad (2.2)$$

or, employing again the stereographic projection,

$$|z\rangle = \sum_{m=-S}^S \binom{2S}{S+m}^{\frac{1}{2}} [\sin(\theta/2)]^{S+m} [\cos(\theta/2)]^{S-m} \times e^{-i(S+m)\phi} |S, m\rangle. \quad (2.3)$$

The system of spin coherent states is complete, but the states are not mutually orthogonal; their overlap is

$$\langle z|z'\rangle = \frac{(1 + z^*z')^{2S}}{[(1 + |z|^2)(1 + |z'|^2)]^S}. \quad (2.4)$$

They allow for a resolution of the unity in the form

$$\int_{\mathbb{C}} d\mu_S(z) |z\rangle\langle z| = \mathbb{1}, \quad (2.5)$$

with the invariant measure given by

$$d\mu_S(z) = \frac{2S + 1}{\pi} \frac{d^2z}{(1 + |z|^2)^2}. \quad (2.6)$$

With this completeness relation one is able to decompose an arbitrary pure state over the coherent states. If we denote $\psi(z^*) = \langle z|\psi\rangle$, by using the basis $\{|S, m\rangle\}$, we define the stellar function $f_{\psi}(z)$ of the state $|\psi\rangle$ as

$$\begin{aligned} f_{\psi}(z) &= (1 + |z|^2)^S \psi(z) = \sum_{m=-S}^S \binom{2S}{S+m}^{\frac{1}{2}} \psi_m z^{S+m} \\ &= \sum_{k=0}^{2S} \binom{2S}{k}^{\frac{1}{2}} \psi_{k-S} z^k, \end{aligned} \quad (2.7)$$

with $\psi_m = \langle S, m|\psi\rangle$, and in the second line we have made the relabeling $S + m \mapsto k$. Interestingly, in this representation the wave function is a polynomial in z of order $r \leq 2S$. In consequence, the roots $z_k \in \mathbb{C}$ of $f_{\psi}(z)$ fully characterize the state. These roots define, via an inverse stereographic map, $2S$ points on the unit sphere \mathcal{S}_2 . This is the Majorana constellation, and each one of these points constitutes one star of the constellation. Note that an $SU(2)$ rotation corresponds to a solid rotation of the constellation; therefore, states with the same constellation, irrespective of their relative orientation, have the same physical properties.

The stellar function is directly related to the Husimi Q function [91,92]:

$$Q_{\psi}(z) = (1 + |z|^2)^{2S} |f_{\psi}(z^*)|^2, \quad (2.8)$$

which clearly shows that the zeros of the Husimi Q_{ψ} function are the complex conjugates of the zeros of f_{ψ} . These can then be observed in the laboratory by measuring where the Husimi function vanishes [93].

Let us examine a few relevant examples to illustrate how these constellations look. The first one is that of a spin coherent state $|z_0\rangle$, whose stellar representation is direct from Eq. (2.4):

$$f_{z_0}(z) = \frac{(1 + z_0z)^{2S}}{(1 + |z_0|^2)^S} \quad (2.9)$$

so it has a single zero at $z = -1/z_0$ with multiplicity $2S$. In consequence, the constellation collapses in this case to a single point diametrically opposed to the maximum z_0 .

Another relevant set of states is that of the so-called NOON states, defined as [94]

$$|\text{NOON}\rangle = \frac{1}{\sqrt{2}} (|S, S\rangle - |S, -S\rangle). \quad (2.10)$$

They are known to have the highest sensitivity for a fixed excitation S to small rotations about the S_z axis [95]. The associated polynomial reads

$$f_{\text{NOON}}(z) = \frac{1}{\sqrt{2}} (z^{2S} - 1). \quad (2.11)$$




FIG. 1. Top: Majorana constellations for, from left to right, a spin coherent state, a NOON state, and a king of quantumness, all of them for the same dimension with $S = 6$. Bottom: Density plots of the corresponding Husimi functions, with a scale that goes from blue (minimum) to red (maximum).

The zeros are thus the $2S$ roots of unity, so the Majorana constellations have $2S$ stars placed around the equator with equal angular separation between each star. A rotation around the S_z axis of angle $\pi/(2S)$ renders the state orthogonal to itself, justifying their optimality.

Finally, we consider kings of quantumness, initially dubbed anticoherent states [96]. In a sense they are the opposite of spin coherent states: whereas the latter correspond as nearly as possible to a classical spin vector pointing in a given direction, the former *point nowhere*; i.e., the average angular momentum vanishes and the fluctuations up to given order M are isotropic [97]. Their symmetrical Majorana constellations herald their isotropic angular momentum properties [98] and correspond to the vertices of Platonic solids in some particular dimensions. In Fig. 1 we show the constellation associated to the examples aforementioned for the case of spin $S = 6$.

III. STATE MULTIPOLES FROM A CONSTELLATION

So far, we have shown how to compute the constellation when the state is given. In this section, we attack the inverse problem: what information can we extract from a given constellation. To this end, we first recall that every polynomial $P(z)$ of degree n can be represented in terms of its zeros ζ_k using the classical Vieta formulas [86], which can be expressed in the form

$$P(z) = \sum_{k=0}^n a_k z^k = a_n \sum_{k=0}^n (-1)^{n-k} e_{n-k}(\boldsymbol{\zeta}) z^k, \quad (3.1)$$

where $e_j(\boldsymbol{\zeta}) \equiv e_j(\zeta_1, \zeta_2, \dots, \zeta_n)$ are the elementary symmetric polynomials [99] defined as

$$\begin{aligned} e_0(\zeta_1, \zeta_2, \dots, \zeta_n) &= 1, \\ e_1(\zeta_1, \zeta_2, \dots, \zeta_n) &= \sum_{1 \leq j \leq n} \zeta_j, \end{aligned}$$

$$\begin{aligned} e_2(\zeta_1, \zeta_2, \dots, \zeta_n) &= \sum_{1 \leq j < k \leq n} \zeta_j \zeta_k, \\ &\vdots \\ e_n(\zeta_1, \zeta_2, \dots, \zeta_n) &= \zeta_1 \zeta_2 \dots \zeta_n. \end{aligned} \quad (3.2)$$

Using this fundamental result, the Majorana stellar function can be expressed in the compact form

$$f_\psi(z) = \sum_{k=0}^{2S} \mathfrak{f}_k(\boldsymbol{\zeta}) z^k, \quad (3.3)$$

where we have introduced the notation

$$\mathfrak{f}_k(\boldsymbol{\zeta}) = (-1)^{2S-k} \psi_S e_{2S-k}(\boldsymbol{\zeta}), \quad (3.4)$$

and the coefficient ψ_S is fixed by the normalization condition

$$\psi_S(\boldsymbol{\zeta}) = \left(\sum_{k=0}^{2S} \frac{|e_{2S-k}(\boldsymbol{\zeta})|^2}{\binom{2S}{k}} \right)^{-1/2}. \quad (3.5)$$

In this way, the state coefficients are simply related to the coefficients of the Majorana polynomial, and we can calculate them as a function of the stars:

$$\psi_{k-S}(\boldsymbol{\zeta}) = \frac{\mathfrak{f}_k(\boldsymbol{\zeta})}{\sqrt{\binom{2S}{k}}}. \quad (3.6)$$

For many purposes, expanding in the basis $|S, m\rangle$ is not a good choice. Instead, if one considers the associated density matrix $\varrho = |\psi\rangle\langle\psi|$, it proves more convenient to use the irreducible tensors [83]. They are defined as

$$T_{Kq} = \sqrt{\frac{2K+1}{2S+1}} \sum_{m,m'=-S}^S C_{Sm,Kq}^{Sm'} |S, m'\rangle\langle S, m|, \quad (3.7)$$

with $C_{Sm,Kq}^{Sm'}$ being the Clebsch-Gordan coefficients that couple a spin S and an integer spin K ($0 \leq K \leq 2S$) to a total spin S [100]. These tensors constitute an orthonormal basis $\text{Tr}(T_{Kq} T_{K'q'}^\dagger) = \delta_{KK'} \delta_{qq'}$ and have the correct transformation properties under rotations.

The corresponding expansion coefficients $\varrho_{Kq} = \text{Tr}(\varrho T_{Kq}^\dagger)$ are known as state multipoles. Actually, ϱ_{Kq} can be related to the K th powers of the generators. The monopole $\varrho_{00} = 1/\sqrt{2S+1}$ is trivially fixed by normalization; the dipole ϱ_{1q} is the first-order moment of \mathbf{S} and thus corresponds to the classical picture, in which the state is represented by its average value on the Bloch sphere. However, the complete characterization of a state demands the knowledge of the other multipoles that account for higher-order fluctuations. The hermiticity of ϱ imposes the conditions $\varrho_{K-q} = (-1)^q \varrho_{Kq}^*$.

The multipoles can be expressed in terms of the state amplitudes ψ_m , which, in turn, can be computed from the constellation. This allows us to calculate ϱ_{Kq} as a function of the constellation in a very compact form:

$$\varrho_{Kq} = \sqrt{\frac{2K+1}{2S+1}} \sum_{m=-S}^S C_{Sm,Kq}^{S,m+q} \psi_{m+q}^*(\boldsymbol{\zeta}) \psi_m(\boldsymbol{\zeta}). \quad (3.8)$$




FIG. 2. Multipole lengths for the three constellations depicted in Fig. 1, as a function of the order K .

It will prove convenient to characterize the multipoles by their effective length:

$$q_K^2 = \sum_{q=-K}^K |q_{Kq}|^2, \quad (3.9)$$

which gauges the state overlapping with the K th multipole pattern and is unchanged via $SU(2)$ rotations of the state. Note that, for pure states we are considering, $\sum_{K=0}^{2S} q_K^2 = 1$. These coefficients q_K^2 have been used as measures of localization [101]; as quantifiers of quantumness [76], which is useful for applications such as rotation sensing [43]; and to quantify mode-decomposition-independent entanglement properties [14].

IV. EXAMPLES

To check how the method works we consider the constellations depicted in Fig. 1, corresponding to the states worked out in Sec. II, namely, coherent, NOON, and king of quantumness for $S = 6$. In Fig. 2 we represent the corresponding q_K^2 as a function of the order K (excluding the monopole, as it is always trivial). Similar patterns emerge for other S . For completeness, in Fig. 3 we also plot the distribution of q_K^2 as a function of the order K and the spin S for the same three states.

In the Bloch sphere, constellations having their points arranged as symmetrically as possible are the most quantum, whereas the opposite occurs for coherent states. As we can appreciate, the coherent state conveys all the relevant information in the lowest-order multipoles, which is in agreement with its classical character. Only for high values of S , one can see a tiny contribution of higher-order multipoles. In the limit of large S , we can fit a Gaussian distribution to these multipoles. Since the distributions are unchanged by rotations of the Bloch sphere, we can choose the coherent state with $z = 0$ to find

$$q_{K,\text{coh}}^2 = \frac{(2K+1)(2S)!^2}{(2S-K)!(2S+K+1)!}. \quad (4.1)$$






FIG. 3. Multipole lengths as a function of the order K and the system spin S for (upper panel) spin coherent states, (middle panel) NOON states, and (lower panel) kings of quantumness. Since the value of the monopole depends on the dimension, we normalized the rest of the multipole lengths to add up to 1. This allows us to compare their values for different S .

The maximum of this distribution is at $K_{\text{max}} = \sqrt{S+1/2} - 1/2$. In the limit of large S , this peaks at $q_{K_{\text{max}}}^2 \approx 1/\sqrt{Se}$, with a variance $S/2$.

The NOON state has only even multipoles which, surprisingly, are identical with the corresponding ones from the coherent state. We thus could regard the NOON state as a coherent state wherein the information from the odd multipoles has been transferred to the highest-order one. In the limit of high S only that multipole is relevant (see Fig. 3),




FIG. 4. Maximum multipole length for each order K , selected from randomly generated states (blue bars). Different states maximize the multipoles for different values of K , so the dotted curves represent the multipoles of the states maximizing a certain region, except for the orange dots, which denote the distribution of the coherent state. The black dot above the last multipole is the value corresponding to the NOON state. Note that the distribution for the coherent state and the state maximizing the last multipole are shown in front of the maximum values, and the rest are vertically displaced for clarity.

confirming that NOON states are more quantum for higher S . The significant contribution of the NOON state to the final multipole raises the question of the maximum value that a state can achieve in its highest-order multipole. To explore this question, we numerically generated 6×10^4 random constellations for various values of S : none of them exhibited a larger highest-order multipole than the NOON state. This brings the question of the maximum value for the K th multipole length.

The results are shown in Fig. 4 for $S = 60$. Unexpectedly, it appears that the coherent state is the unique state with the maximal multipole ϱ_K^2 for all values of K up to a certain point. After this, another state maximizes the K th multipole for a different region of larger K values and so on. The multipolar distribution of these states is bell shaped near the multipole it maximizes and exhibits a wiggling behavior at the first multipoles. We can also observe how, as K increases, this wiggling evolves to a distribution very similar to that of the NOON state. This supports the hypothesis that the NOON state is the one with the highest contribution to its final multipole and allows a discrete transition between states maximizing low-order multipoles (coherent) and those maximizing high-order ones (NOON).

The king of quantumness shows the absence of lower-order multipoles, which is the origin of the isotropic behavior of their higher-order fluctuations. However, the significance of the strength of the first nonzero multipole is not clear.

We can also gain some general intuition from our calculations. If one has a constellation and makes a change, such as adding, removing, or moving a star, what happens to the multipoles? Upon the addition of a new star ζ_{new} to the constellation, for example, the elementary symmetric polynomials change as $e_k \mapsto \zeta_{\text{new}} e_{k-1} + e_k$ for $k > 0$ and the rest of the calculations proceed as above. Supposing further that the original constellation corresponded to a coherent state at $z = 0$, the addition of a star at ζ_{new} makes the multipoles

transform to

$$\varrho_K^2 = c(K, S) \left[4S^2 + 4S|\zeta_{\text{new}}|^2 + \frac{(K^2 + K - 2S)^2}{4S^2} |\zeta_{\text{new}}|^4 \right], \quad (4.2)$$

with

$$c(K, S) = \frac{(2K+1)(2S-1)!^2 |\psi_S|^4}{(2S-K)!(2S+K+1)!}. \quad (4.3)$$

V. DISCUSSION

We explore in this section some nontrivial consequences of our method. A first direct result is to compute Stokes operators [102] from the constellation. Choosing the S_z operator, for example, all of its moments may now be computed as

$$\langle S_z^n \rangle = |\mathfrak{f}_{2S}(\boldsymbol{\zeta})|^2 \sum_{m=0}^{2S} (m-S)^n \binom{2S}{m}^{-1} |e_{2S-m}(\boldsymbol{\zeta})|^2; \quad (5.1)$$

all other moments may be found by rigidly rotating the constellation to highlight any other operator axis. Knowledge of the Stokes vector $\mathbf{S} = (\langle S_x \rangle, \langle S_y \rangle, \langle S_z \rangle)^\top$ (the superscript \top denoting the transpose) and $2S - 1$ of the stars provides an overcomplete set of three equations for finding the location of the remaining star. With two stars' locations unknown, the covariances between the Stokes parameters provide sufficient information to determine the four unknown angular coordinate parameters, and so on for higher-order moments and determining the unknown locations of more stars. The inversion process involves nonlinear functions and tends to require numerical solutions.

The results may also be used to make contact with spherical t designs, which are sets of points on the sphere that may be used for averaging polynomial functions over the entire sphere [58,59]. For a set of points to be a one-design, their vectors must sum to the zero vector. In terms of the stereographic projection, this constrains the points to obey

$$\sum_j \frac{\zeta_j}{1 + |\zeta_j|^2} = 0, \quad \sum_j \frac{1 - |\zeta_j|^2}{1 + |\zeta_j|^2} = 0. \quad (5.2)$$

More connections to designs with $t > 1$ are the subject of future study.

We can further explore the significance of the elementary symmetric polynomials, as they have not yet been used to our knowledge for studying our quantum states. Consider that a Majorana constellation can always be rotated such that one of the stars is at the north pole with $\theta = 0$. This makes the highest-order polynomial e_{2S} vanish. Then, the vanishing of the next polynomial e_{2S-1} implies that a second star is also at the north pole. With each subsequent polynomial that vanishes, another star is added to the north pole, up to the condition that $e_j = 0 \forall j > 0$ implies that all $2S$ stars are at the north pole and the state is a spin coherent state. The maximal degeneracy of any star in the constellation is then equal to the maximum number of consecutive highest-order symmetric polynomials that vanish, where the latter is maximized over rigid rotations of the stars that perform a known transformation of the roots.




FIG. 5. Two visualizations of how a Majorana constellation can smoothly evolve between classical (coherent) and quantum (NOON) states. On the left, the stars are always at equal polar angles to each other and are equally spaced from each other, while the circle they are spaced on transitions between the north pole (coherent) and the equator (NOON). On the right, the stars spread or coalesce around the equator, with equal azimuthal angles between each star and the subsequent star; the final star goes around the equator to come closer to (NOON) or farther from (coherent) wrapping around to reach the initial star from the other side.

Specifying which polynomials $e_j(\zeta)$ vanish significantly constrains a state. In quantifying quantumness, the vanishing of low-order multipoles is a sign of nonclassicality, while here the vanishing of all higher-order polynomials seems to imply an increase in classicality. We are led to consider an interesting case: what if all of the *lower*-order polynomials vanish, such that the state retains $e_{2S} \neq 0$ while $e_j(\zeta) = 0 \forall 0 < j < 2S$? This includes, for example, the NOON states, which are considered highly quantum by many measures. Since NOON states have all of their stars equally spaced around the equator, the roots all obey $|\zeta_j| = |\zeta_k|$ and are directly equal to roots of unity $\zeta_j = \exp(i\pi j/S)$. Summing all $2S$ such roots gives $e_1(\zeta) = \sum_{j=1}^{2S} \exp(i\pi j/S) = 0$; summing all $\binom{2S}{2}$ products of such roots gives

$$\begin{aligned} e_2(\zeta) &= \sum_{j < k}^{2S} \exp[i\pi(j+k)/S] \\ &= \frac{1}{2} \sum_{j \neq k}^{2S} \exp[i\pi(j+k)/S] = 0, \end{aligned} \quad (5.3)$$

and so on for all higher-order $e_j(\zeta)$ other than

$$e_{2S}(\zeta) = \exp\left(i\pi \sum_{j=1}^{2S} \frac{j}{S}\right) = (-1)^{2S+1} \neq 0. \quad (5.4)$$

We can visually and mathematically explore the transition between NOON states and coherent states by the migration of stars from the equator to the north pole and by the vanishing of $e_{2S}(\zeta)$, respectively. Consider the transition as sketched in Fig. 5(a), where the $2S$ points in the constellation are equally spaced around a circle other than the equator, sharing the polar angle θ . The symmetric polynomials take the same form as before, with $e_j(\zeta)$ multiplied by $\tan^j(\theta/2)$, so all of the lowest polynomials vanish other than

$$e_{2S} = (-1)^{2S+1} \tan^{2S}(\theta/2). \quad (5.5)$$




FIG. 6. Evolution of the multipole lengths corresponding to Fig. 5(a). The position of the stars in the figure is for illustrative purposes only, as the transition occurs when the stars are very close to the equator.

This codifies the gradual progression: as the circle on which the stars lie gets smaller and smaller, from a NOON to a coherent state, the one nonzero polynomial (other than $e_0 = 1$) goes to zero as $\tan^{2S}(\theta/2)$. The magnitude of this highest-order polynomial, after a rigid rotation that makes the rest of the polynomials vanish, directly encodes the quantumness of the state. Note that it is the deviation of the magnitude of e_{2S} from unity that matters: there is a coherent state at $\theta = 0$ and π .

We can also obtain exact expressions for the multipoles as a function of θ :

$$Q_K^2 = Q_{K,\text{coh}}^2 \left[\frac{\tan^{4S}(\theta/2) - 1}{\tan^{4S}(\theta/2) + 1} \right]^2, \quad K = 1, 3, \dots, 2S - 1, \quad (5.6)$$

whereas $Q_K^2 = Q_{K,\text{coh}}^2$ for $K = 2, 4, \dots, 2S - 2$. This transition from coherent to NOON is very abrupt, even for low values of S , due to the power of $4S$ in the tangent. The multipoles look much like a coherent state up until the stars are very close to the equator. This shows the vulnerability of the NOON state to a small displacement of its stars.

As previously argued, the NOON state maximizes the last multipole. With the help of the previous expressions we can obtain this value:

$$Q_{2S,\text{NOON}}^2 = \begin{cases} \frac{1}{2} + \binom{4S}{2S}^{-1} & S = 1, 2, \dots \\ \frac{1}{2} & S = \frac{1}{2}, \frac{3}{2}, \dots \end{cases}, \quad (5.7)$$

which tends to $1/2$ as S becomes large. Note that this does not contradict the results in Fig. 3, for the multipoles are normalized therein.

We can observe how the multipole lengths change in the aforementioned transition between coherent and NOON states in Fig. 6. This transition can be enacted by evolution under the highly nontrivial Hamiltonian

$$H_{\text{NOON} \leftrightarrow \text{coh}} = -i|S, S\rangle\langle S, -S| + i|S, -S\rangle\langle S, S| \quad (5.8)$$

that manifestly breaks degeneracies between Majorana stars.

Further consider the alternative transition as in Fig. 5(b), where the stars spread around the equator ($\theta = \pi/2$) smoothly

between NOON- and coherent-state configurations. Now we consider the coherent state to have $\zeta = 1$, such that its polynomials take the form $e_j(\zeta) = \binom{2S}{j}$. We choose the spreading rate such that all stars arrive in place at the same time, with $\zeta_j(t) = \exp(i\pi t j/S)$ and $t \in (0, 1)$ with $t = 0$ and 1 corresponding to coherent and NOON states, respectively. Then

$$\begin{aligned} e_1[\zeta(t)] &= \exp\left[i\pi \frac{(2S+1)}{2S}t\right] \frac{\sin(\pi t)}{\sin\left(\frac{\pi t}{2S}\right)}, \\ e_2[\zeta(t)] &= \frac{1}{2!}\{e_1[\zeta(t)]^2 - e_1[\zeta(2t)]\}, \\ e_3[\zeta(t)] &= \frac{1}{3!}\{e_1[\zeta(t)]^3 - 3e_1[\zeta(t)]e_1[\zeta(2t)] + 2e_1[\zeta(3t)]\}, \end{aligned} \quad (5.9)$$

and so on. For small t , these equal

$$e_j[\zeta(t)] = \binom{2S}{j} - \binom{2S}{j+1} \pi^2 \frac{2S+1}{4S^2} t^2 + \mathcal{O}(t^4). \quad (5.10)$$

The Hamiltonian required to enact this transition is much more complicated, as the stars all rotate around the z axis but no Hamiltonian of the form S_z^n can do anything other than rotate the NOON state.

Finally, consider the symmetric transition where the stars spread in both directions along the equator, with pairs moving at equal speeds. For example, with $2S$ odd, we can set $\zeta_0 = 1$, $\zeta_j(t) = \exp(i\pi t j/S)$ for $0 < j \leq S - 1/2$, and relabel the remainder as $\zeta_{-j}(t) = \exp(-i\pi t j/S)$ for $0 < j \leq S - 1/2$. The spacings between all of the stars are the same as when all of the stars traveled in the same direction, merely offset by a relative phase, which symmetrizes the expressions to

$$\begin{aligned} e_1[\zeta(t)] &= \frac{\sin(\pi t)}{\sin\left(\frac{\pi t}{2S}\right)}, \\ e_2[\zeta(t)] &= \frac{1}{2!}\{e_1[\zeta(t)]^2 - e_1[\zeta(2t)]\}, \end{aligned} \quad (5.11)$$

and so on. We can thus use the symmetric polynomials to observe the transitions between highly quantum and highly

classical states, in accordance with the geometric picture of the Majorana constellation.

VI. CONCLUDING REMARKS

The Majorana representation provides a valuable geometric tool to characterize quantum states with $SU(2)$ symmetry. The visualization of quantum states as a constellation in the unit sphere is finding new applications in quantum information science. On the other hand, the multipolar expansion is especially germane to capture the state invariant properties and, in addition, the resulting multipoles can be measured in the laboratory.

We have extensively explored how to go from Majorana constellations to multipolar distributions and back, illustrating this through a variety of examples. For instance, we have elucidated the changes in the multipolar distribution when the Majorana constellation transitions from a classical (concentrated) to a highly quantum state (spread out). Additionally, we have delved into the consequences of adding new stars to the constellation and examined states with the greatest contribution to the higher-order multipole, among many other aspects. Interestingly, the formalism can be manifestly extended to other symmetries [103]: this is more than an academic curiosity, and work in this direction is ongoing.

ACKNOWLEDGMENTS

We are indebted to G. Björk, P. de la Hoz, M. Grassl, L. Villanueva, and Ł. Rudnicki for discussions. A.Z.G. acknowledges that the NRC headquarters is located on the traditional unceded territory of the Algonquin Anishinaabe and Mohawk people. We acknowledge financial support from the European Union QUANTERA program (project ApresSF) and the Spanish Research Agency (Grant No. PID2021-127781NB-I00). A.Z.G. acknowledges funding from the Natural Sciences and Engineering Research Council PDF program.

-
- [1] K. Eckert, J. Schliemann, D. Bruß, and M. Lewenstein, Quantum correlations in systems of indistinguishable particles, *Ann. Phys.* **299**, 88 (2002).
- [2] Y. Kawaguchi and M. Ueda, Spinor Bose-Einstein condensates, *Phys. Rep.* **520**, 253 (2012).
- [3] D. M. Stamper-Kurn and M. Ueda, Spinor Bose gases: Symmetries, magnetism, and quantum dynamics, *Rev. Mod. Phys.* **85**, 1191 (2013).
- [4] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Observation of Bose-Einstein condensation in a dilute atomic vapor, *Science* **269**, 198 (1995).
- [5] K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Bose-Einstein condensation in a gas of sodium atoms, *Phys. Rev. Lett.* **75**, 3969 (1995).
- [6] M. S. Chang, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M. Fortier, W. Zhang, L. You, and M. S. Chapman, Observation of spinor dynamics in optically trapped ^{87}Rb Bose-Einstein condensates, *Phys. Rev. Lett.* **92**, 140403 (2004).
- [7] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, Bose-Einstein condensation of chromium, *Phys. Rev. Lett.* **94**, 160401 (2005).
- [8] Q. Beaufils, R. Chicireanu, T. Zanon, B. Laburthe-Tolra, E. Maréchal, L. Vernac, J. C. Keller, and O. Gorceix, All-optical production of chromium Bose-Einstein condensates, *Phys. Rev. A* **77**, 061601(R) (2008).
- [9] A. R. Usha Devi, R. Prabhu, and A. K. Rajagopal, Characterizing multiparticle entanglement in symmetric n -qubit states via negativity of covariance matrices, *Phys. Rev. Lett.* **98**, 060501 (2007).
- [10] T. Bastin, S. Krins, P. Mathonet, M. Godefroid, L. Lamata, and E. Solano, Operational families of entanglement classes for symmetric n -qubit states, *Phys. Rev. Lett.* **103**, 070503 (2009).

- [11] J. Martin, O. Giraud, P. A. Braun, D. Braun, and T. Bastin, Multiqubit symmetric states with high geometric entanglement, *Phys. Rev. A* **81**, 062347 (2010).
- [12] M. Aulbach, Classification of entanglement in symmetric states, *Int. J. Quantum Inform.* **10**, 1230004 (2012).
- [13] A. Mandilara, T. Coudreau, A. Keller, and P. Milman, Entanglement classification of pure symmetric states via spin coherent states, *Phys. Rev. A* **90**, 050302(R) (2014).
- [14] A. Z. Goldberg, M. Grassl, G. Leuchs, and L. L. Sánchez-Soto, Quantumness beyond entanglement: The case of symmetric states, *Phys. Rev. A* **105**, 022433 (2022).
- [15] E. Majorana, Atomi orientati in campo magnetico variabile, *Nuovo Cim.* **9**, 43 (1932).
- [16] F. Bloch and I. I. Rabi, Atoms in variable magnetic fields, *Rev. Mod. Phys.* **17**, 237 (1945).
- [17] H. Salwen, Resonance transitions in molecular beam experiments. I. General theory of transitions in a rotating magnetic field, *Phys. Rev.* **99**, 1274 (1955).
- [18] A. Meckler, Majorana formula, *Phys. Rev.* **111**, 1447 (1958).
- [19] J. Schwinger, The Majorana formula, *Trans. NY Acad. Sci.* **38**, 170 (1977).
- [20] R. Penrose, A spinor approach to general relativity, *Ann. Phys.* **10**, 171 (1960).
- [21] R. Penrose, *The Emperor's New Mind* (Oxford University, New York, 1989).
- [22] H. Bacry, *Group Theory and Constellations* (Publibook, Paris, 2004).
- [23] J. H. Hannay, The Majorana representation of polarization, and the Berry phase of light, *J. Mod. Opt.* **45**, 1001 (1998).
- [24] G. Björk, A. B. Klimov, P. de la Hoz, M. Grassl, G. Leuchs, and L. L. Sánchez-Soto, Extremal quantum states and their Majorana constellations, *Phys. Rev. A* **92**, 031801(R) (2015).
- [25] A. Z. Goldberg, Quantum theory of polarimetry: From quantum operations to Mueller matrices, *Phys. Rev. Res.* **2**, 023038 (2020).
- [26] A. Z. Goldberg, P. de la Hoz, G. Björk, A. B. Klimov, M. Grassl, G. Leuchs, and L. L. Sánchez-Soto, Quantum concepts in optical polarization, *Adv. Opt. Photon.* **13**, 1 (2021).
- [27] A. Z. Goldberg, Quantum polarimetry, *Prog. Opt.* **67**, 185 (2022).
- [28] R. Barnett, D. Podolsky, and G. Refael, Geometrical approach to hydrodynamics and low-energy excitations of spinor condensates, *Phys. Rev. B* **80**, 024420 (2009).
- [29] A. Lamacraft, Low-energy dynamics of spinor condensates, *Phys. Rev. B* **81**, 184526 (2010).
- [30] B. Lian, T.-L. Ho, and H. Zhai, Searching for non-Abelian phases in the Bose-Einstein condensate of dysprosium, *Phys. Rev. A* **85**, 051606(R) (2012).
- [31] X. Cui, B. Lian, T.-L. Ho, B. L. Lev, and H. Zhai, Synthetic gauge field with highly magnetic lanthanide atoms, *Phys. Rev. A* **88**, 011601(R) (2013).
- [32] H. Mäkelä and A. Messina, N -qubit states as points on the Bloch sphere, *Phys. Scr.* **2010**, 014054 (2010).
- [33] P. Ribeiro and R. Mosseri, Entanglement in the symmetric sector of n qubits, *Phys. Rev. Lett.* **106**, 180502 (2011).
- [34] A. R. U. Devi, Sudha, and A. K. Rajagopal, Majorana representation of symmetric multiqubit states, *Quantum Inf. Process.* **11**, 685 (2012).
- [35] C. Chryssomalakos, L. Hanotel, E. Guzmán-González, D. Braun, E. Serrano-Ensástiga, and K. Życzkowski, Symmetric multiqudit states: Stars, entanglement, and roto-sensors, *Phys. Rev. A* **104**, 012407 (2021).
- [36] C.-F. Kam and R.-B. Liu, Three-tangle of a general three-qubit state in the representation of majorana stars, *Phys. Rev. A* **101**, 032318 (2020).
- [37] X. Lu, L.-Y. Zhan, J. Chen, H.-D. Liu, L.-B. Fu, and X.-G. Wang, Many-body anticommutator and its applications in the normalization of completely symmetric states with Majorana's stellar representations, *Phys. Rev. A* **104**, 012203 (2021).
- [38] A. Burchardt, J. Czartowski, and K. Życzkowski, Entanglement in highly symmetric multipartite quantum states, *Phys. Rev. A* **104**, 022426 (2021).
- [39] F. Bouchard, P. de la Hoz, G. Björk, R. W. Boyd, M. Grassl, Z. Hradil, E. Karimi, A. B. Klimov, G. Leuchs, J. Řeháček, and L. L. Sánchez-Soto, Quantum metrology at the limit with extremal Majorana constellations, *Optica* **4**, 1429 (2017).
- [40] C. Chryssomalakos and H. Hernández-Coronado, Optimal quantum roto-sensors, *Phys. Rev. A* **95**, 052125 (2017).
- [41] A. Z. Goldberg and D. F. V. James, Quantum-limited Euler angle measurements using anticoherent states, *Phys. Rev. A* **98**, 032113 (2018).
- [42] J. Martin, S. Weigert, and O. Giraud, Optimal detection of rotations about unknown axes by coherent and anticoherent states, *Quantum* **4**, 285 (2020).
- [43] A. Z. Goldberg, A. B. Klimov, G. Leuchs, and L. L. Sánchez-Soto, Rotation sensing at the ultimate limit, *J. Phys. Photonics* **3**, 022008 (2021).
- [44] J. H. Hannay, The Berry phase for spin in the Majorana representation, *J. Phys. A* **31**, L53 (1998).
- [45] P. Bruno, Quantum geometric phase in Majorana's stellar representation: Mapping onto a many-body Aharonov-Bohm phase, *Phys. Rev. Lett.* **108**, 240402 (2012).
- [46] H. D. Liu and L. B. Fu, Representation of Berry phase by the trajectories of Majorana stars, *Phys. Rev. Lett.* **113**, 240403 (2014).
- [47] C. Yang, H. Guo, L.-B. Fu, and S. Chen, Characterization of symmetry-protected topological phases in polymerized models by trajectories of Majorana stars, *Phys. Rev. B* **91**, 125132 (2015).
- [48] H. D. Liu and L. B. Fu, Berry phase and quantum entanglement in Majorana's stellar representation, *Phys. Rev. A* **94**, 022123 (2016).
- [49] F. Yao, D. Li, H. Liu, L. Fu, and X. Wang, The exact curve equation for Majorana stars, *Sci. Rep.* **7**, 15558 (2017).
- [50] C.-F. Kam and R.-B. Liu, Berry phases of higher spins due to internal geometry of Majorana constellation and relation to quantum entanglement, *New J. Phys.* **23**, 073020 (2021).
- [51] V. Mittal, K. S. Akhilesh, and S. K. Goyal, Geometric decomposition of geodesics and null-phase curves using Majorana star representation, *Phys. Rev. A* **105**, 052219 (2022).
- [52] J. Bartlett, H. Hu, and E. Zhao, Illuminating the bulk-boundary correspondence of a non-Hermitian stub lattice with Majorana stars, *Phys. Rev. B* **104**, 195131 (2021).
- [53] P. Ribeiro, J. Vidal, and R. Mosseri, Thermodynamical limit of the Lipkin-Meshkov-Glick model, *Phys. Rev. Lett.* **99**, 050402 (2007).
- [54] P. Ribeiro, J. Vidal, and R. Mosseri, Exact spectrum of the Lipkin-Meshkov-Glick model in the thermodynamic limit and finite-size corrections, *Phys. Rev. E* **78**, 021106 (2008).

- [55] D. J. H. Markham, Entanglement and symmetry in permutation-symmetric states, *Phys. Rev. A* **83**, 042332 (2011).
- [56] Z. Wang and D. Markham, Nonlocality of symmetric states, *Phys. Rev. Lett.* **108**, 210407 (2012).
- [57] W. Ganczarek, M. Kuś, and K. Życzkowski, Barycentric measure of quantum entanglement, *Phys. Rev. A* **85**, 032314 (2012).
- [58] P. Delsarte, J. M. Goethals, and J. J. Seidel, Spherical codes and designs, *Geom. Dedicata* **6**, 363 (1977).
- [59] R. H. Hardin and N. J. A. Sloane, New spherical 4-designs, *Discrete Math.* **106-107**, 255 (1992).
- [60] J. J. Thomson, On the structure of the atom: An investigation of the stability and periods of oscillation of a number of corpuscles arranged at equal intervals around the circumference of a circle; with application of the results to the theory of atomic structure, *Philos. Mag. Ser. 7*, 37265 (1904).
- [61] T. H. Melnyk, O. Knop, and W. R. Smith, Extremal arrangements of points and unit charges on a sphere: Equilibrium configurations revisited, *Can. J. Chem.* **55**, 1745 (1977).
- [62] N. Ashby and W. E. Brittin, Thomson's problem, *Am. J. Phys.* **54**, 776 (1986).
- [63] T. Erber and G. M. Hockney, Equilibrium configurations of n equal charges on a sphere, *J. Phys. A* **24**, L1369 (1991).
- [64] J. R. Edmundson, The distribution of point charges on the surface of a sphere, *Acta Cryst.* **A48**, 46 (1992).
- [65] A. Baecklund and I. Bengtsson, Four remarks on spin coherent states, *Phys. Scr.* **2014**, 014012 (2014).
- [66] P. M. L. Tammes, On the origin of number and arrangement of places of exit on the surface of pollen grains, *Recueil Trav. Bot. Néerl.* **27**, 1 (1930).
- [67] L. Fejes Tóth, *Lagerungen in der Ebene, auf der Kugel und in Raum* (Springer-Verlag, Berlin, 1953).
- [68] J. Leech, Equilibrium of sets of particles on a sphere, *Math. Gaz.* **41**, 81 (1957).
- [69] O. Giraud, P. Braun, and D. Braun, Quantifying quantumness and the quest for queens of quantumness, *New J. Phys.* **12**, 063005 (2010).
- [70] G. Björk, M. Grassl, P. de la Hoz, G. Leuchs, and L. L. Sánchez-Soto, Stars of the quantum universe: Extremal constellations on the Poincaré sphere, *Phys. Scr.* **90**, 108008 (2015).
- [71] M. Aulbach, D. Markham, and M. Muraio, The maximally entangled symmetric state in terms of the geometric measure, *New J. Phys.* **12**, 073025 (2010).
- [72] L. Arnaud and N. J. Cerf, Exploring pure quantum states with maximally mixed reductions, *Phys. Rev. A* **87**, 012319 (2013).
- [73] D. Goyeneche and K. Życzkowski, Genuinely multipartite entangled states and orthogonal arrays, *Phys. Rev. A* **90**, 022316 (2014).
- [74] M.-S. Li and Y.-L. Wang, k -uniform quantum states arising from orthogonal arrays, *Phys. Rev. A* **99**, 042332 (2019).
- [75] E. H. Lieb and J. P. Solovej, Proof of an entropy conjecture for Bloch coherent spin states and its generalizations, *Acta Math.* **212**, 379 (2014).
- [76] A. Z. Goldberg, A. B. Klimov, M. Grassl, G. Leuchs, and L. L. Sánchez-Soto, Extremal quantum states, *AVS Quantum Science* **2**, 044701 (2020).
- [77] L. L. Whyte, Unique arrangements of points on a sphere, *Am. Math. Monthly* **59**, 606 (1952).
- [78] E. B. Saff and A. B. J. Kuijlaars, Distributing many points on a sphere, *Math. Intell.* **19**, 5 (1997).
- [79] A. Katanforoush and M. Shahshahani, Distributing points on the sphere, I, *Exp. Math.* **12**, 199 (2003).
- [80] J. S. Brauchart and P. J. Grabner, Distributing many points on spheres: Minimal energy and designs, *J. Complexity* **31**, 293 (2015).
- [81] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (Wiley, New York, 1999).
- [82] U. Fano and G. Racah, *Irreducible Tensorial Sets* (Academic, New York, 1959).
- [83] K. Blum, *Density Matrix Theory and Applications* (Plenum, New York, 1981).
- [84] G. Björk, J. Söderholm, Y. S. Kim, Y. S. Ra, H. T. Lim, C. Kothe, Y. H. Kim, L. L. Sánchez-Soto, and A. B. Klimov, Central-moment description of polarization for quantum states of light, *Phys. Rev. A* **85**, 053835 (2012).
- [85] A. Z. Goldberg, A. B. Klimov, H. deGuise, G. Leuchs, G. S. Agarwal, and L. L. Sánchez-Soto, From polarization multipoles to higher-order coherences, *Opt. Lett.* **47**, 477 (2022).
- [86] H. G. Funkhouser, A short account of the history of symmetric functions of roots of equations, *Am. Math. Monthly* **37**, 357 (1930).
- [87] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States* (Cambridge University, Cambridge, England, 2006).
- [88] A. Perelomov, *Generalized Coherent States and their Applications* (Springer-Verlag, Berlin, 1986).
- [89] J.-P. Gazeau, *Coherent States in Quantum Physics* (Wiley, New York, 2009).
- [90] H. S. M. Coxeter, *Introduction to Geometry* (Wiley, New York, 1969).
- [91] K. Husimi, Some formal properties of the density matrix, *Proc. Phys. Math. Soc. Jpn.* **22**, 264 (1940).
- [92] Y. Kano, A new phase-space distribution function in the statistical theory of the electromagnetic field, *J. Math. Phys.* **6**, 1913 (1965).
- [93] C. R. Müller, C. Peuntinger, T. Dirmeier, I. Khan, U. Vogl, C. Marquardt, G. Leuchs, L. L. Sánchez-Soto, Y. S. Teo, Z. Hradil, and J. Řeháček, Evading Vacuum Noise: Wigner Projections or Husimi Samples? *Phys. Rev. Lett.* **117**, 070801 (2016).
- [94] J. P. Dowling, Quantum optical metrology: The lowdown on high-N00N states, *Contemp. Phys.* **49**, 125 (2008).
- [95] J. J. Bollinger, W. M. Itano, D. J. Wineland, and D. J. Heinzen, Optimal frequency measurements with maximally correlated states, *Phys. Rev. A* **54**, R4649(R) (1996).
- [96] J. Zimba, "Anticoherent" spin states via the Majorana representation, *Electron. J. Theor. Phys.* **3**, 143 (2006).
- [97] P. de la Hoz, A. B. Klimov, G. Björk, Y. H. Kim, C. Müller, C. Marquardt, G. Leuchs, and L. L. Sánchez-Soto, Multipolar hierarchy of efficient quantum polarization measures, *Phys. Rev. A* **88**, 063803 (2013).
- [98] <http://polarization.markus-grassl.de>.
- [99] B. van der Waerden, *Algebra* (Springer-Verlag, Berlin, 1991), Vol. 1.
- [100] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonski, *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).
- [101] M. J. W. Hall, Universal geometric approach to uncertainty, entropy, and information, *Phys. Rev. A* **59**, 2602 (1999).

- [102] A. Luis and L. L. Sánchez-Soto, Quantum phase difference, phase measurements and Stokes operators, [Prog. Opt.](#) **41**, 421 (2000).
- [103] N. Fabre, A. B. Klimov, R. Murenzi, J.-P. Gazeau, and L. L. Sánchez-Soto, Majorana stellar representation of twisted photons, [Phys. Rev. Res.](#) **5**, L032006 (2023).