NRC Publications Archive Archives des publications du CNRC

Chemical constituents of the physodes of brown algae. Characterization by ^1H and ^{13}C nuclear magnetic resonance spectroscopy of oligomers of phloroglucinol from *Fucus vesiculosus* (L.)

Craigie, James S.; McInnes, A. Gavin; Ragan, Mark A.; Walter, John A.

This publication could be one of several versions: author's original, accepted manuscript or the publisher's version. /
La version de cette publication peut être l'une des suivantes : la version prépublication de l'auteur, la version
acceptée du manuscrit ou la version de l'éditeur.

For the publisher's version, please access the DOI link below. / Pour consulter la version de l'éditeur, utilisez le lien
DOI ci-dessous.

Publisher's version / Version de l'éditeur:

<https://doi.org/10.1139/v77-219>

Canadian Journal of Chemistry, 55, 9, pp. 1575-1582, 1977-05

NRC Publications Record / Notice d'Archives des publications de CNRC:

<https://nrc-publications.canada.ca/eng/view/object/?id=c4349367-9372-499d-8d4a-2a2a03b594e>;

<https://publications-cnrc.canada.ca/fra/voir/objet/?id=c4349367-9372-499d-8d4a-2a2a03b594ea>

Access and use of this website and the material on it are subject to the Terms and Conditions set forth at

<https://nrc-publications.canada.ca/eng/copyright>

READ THESE TERMS AND CONDITIONS CAREFULLY BEFORE USING THIS WEBSITE.

L'accès à ce site Web et l'utilisation de son contenu sont assujettis aux conditions présentées dans le site

<https://publications-cnrc.canada.ca/fra/droits>

LISEZ CES CONDITIONS ATTENTIVEMENT AVANT D'UTILISER CE SITE WEB.

Questions? Contact the NRC Publications Archive team at

PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca. If you wish to email the authors directly, please see the
first page of the publication for their contact information.

Vous avez des questions? Nous pouvons vous aider. Pour communiquer directement avec un auteur, consultez la
première page de la revue dans laquelle son article a été publié afin de trouver ses coordonnées. Si vous n'arrivez
pas à les repérer, communiquez avec nous à PublicationsArchive-ArchivesPublications@nrc-cnrc.gc.ca.

Chemical constituents of the physodes of brown algae. Characterization by ^1H and ^{13}C nuclear magnetic resonance spectroscopy of oligomers of phloroglucinol from *Fucus vesiculosus* (L.)¹

JAMES S. CRAIGIE, A. GAVIN MCINNES, MARK A. RAGAN,²
AND JOHN A. WALTER

Atlantic Regional Laboratory, National Research Council of Canada, Halifax, N.S., Canada B3H 3Z1

Received December 22, 1976

JAMES S. CRAIGIE, A. GAVIN MCINNES, MARK A. RAGAN, and JOHN A. WALTER. Can. J. Chem. **55**, 1575 (1977).

Alcoholic extracts of *Fucus vesiculosus* contain small quantities of low molecular weight polyphenols derived from phloroglucinol and 2,2',4,4',6,6'-hexahydroxybiphenyl. ^1H and ^{13}C nmr were used to identify two of these as 4-(2'',4'',6''-trihydroxyphenoxy)-2,2',4',6,6'-pentahydroxybiphenyl and 4-(2''-(2''',4''',6'''-trihydroxyphenoxy)-4'',6''-dihydroxyphenoxy)-2,2',4',6,6'-pentahydroxybiphenyl.

JAMES S. CRAIGIE, A. GAVIN MCINNES, MARK A. RAGAN et JOHN A. WALTER. Can. J. Chem. **55**, 1575 (1977).

Les produits obtenus par extractions alcooliques du *Fucus vesiculosus* contiennent des petites quantités de polyphénols de bas poids moléculaires dérivés du phloroglucinol et de l'hexahydroxy-2,2',4,4',6,6' biphenyle. On a utilisé la rmn ^1H et ^{13}C pour en identifier deux qui sont le (trihydroxy-2'',4'',6'' phénoxy)-4 pentahydroxy-2,2',4',6,6' biphenyle et le [(trihydroxy-2''',4''',6''' phénoxy)-2'' dihydroxy-4'',6'' phénoxy]-4 pentahydroxy-2,2',4',6,6' biphenyle.

[Traduit par le journal]

Introduction

In 1892 it was demonstrated (1) that brown algal subcellular bodies, called physodes, gave a fiery red colour with the Lindt reagent (vanillin-HCl). This distinctive reaction led to the generally accepted belief that brown algae contained phloroglucinol **1a** or related compounds (2).³ Chemical evidence that brown algal cells produce **1a**, however, was lacking until quite recently when it was demonstrated in hydrolysates of the tannins of *Sargassum ringgoldianum* (3), exudates of *Fucus vesiculosus* (4), and in the direct extracts of seventeen algal species (5). In a previous communication (6) we showed that extracts of Nova Scotian *F. vesiculosus* contained **1a** and

2,2',4,4',6,6'-hexahydroxybiphenyl **2a** ('dimer'), other oligomers, and high molecular weight polymers composed of **1a** and **2a**. Glombitza *et al.* (7) have independently identified the peracetylated derivatives of **1a** and **2a** in acetylated

1a R = H^a
1b R = COCH₃^{a,b}

2a R = H^a
2b R = COCH₃^{a,b}

extracts of *F. vesiculosus*, and also claimed the presence of a terphenyl and two quaterphenyls presumably derived from residues of **1a**. However, on the basis of their published data it is

¹NRCC No. 15848.

²Institute of Marine Biochemistry, University of Trondheim, N-7034, Trondheim, Norway.

³In the structures illustrated the superscripts a-f refer to the following: a, numbers in parentheses are δ (TMS) (ppm); numbers not in parentheses are δ_c (TMS or TSP) (ppm); underlined numbers refer to R = CH₃CO; numbers not underlined to R = H; b, resonances for ^{13}C nuclei in CH₃CO— groups have not been individually assigned; c, data from ref. 13; d, data from ref. 8; e, data from ref. 19; ^{13}C resonances were reassigned by comparison with **5b**, using the substituent effect of a CH₃CO₂ group at C-4' (from **9** and **10**, see text); f, data from ref. 17.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National Research Council of Canada on 02/28/17
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

Can. J. Chem. Downloaded from www.nrcresearchpress.com by National University of Singapore on 05/21/15
For personal use only.

by a phenyl substituent (16). Each of the **3b** hydrogens (H-3, H-5) would, therefore, lie between a phenoxy and an acetoxy (δ 2.13, 3H) substituent, like H-2' and H-6' of **7b**. The presence of only two signals with chemical shifts $\delta \geq 2.20$ from acetoxy methyl groups in **3b** excluded the possibility of a 3,4,5-triacetoxyphenoxy substituent as in **7b**, the spectrum of which contains three such signals. In both **5b** and **7b**, a phenoxy substituent shields the protons of adjacent acetoxy groups. This eliminates **6a** and **6b** as possible structures. The combined 1H nmr results favour structure **3a** for the unknown and **3b** for its acetate derivative.

The structures of **3a** and **3b** were finally established by comparing their ^{13}C spectra with those for the model compounds **1a**, **1b**, **2a**, **2b**,

5a, and **5b**. All of the model compounds possessed symmetry elements which resulted in a reduction in the number of signals observed for the aromatic carbons. Thus the ^{13}C spectra of **1a** and **1b** contained only two resonances for such carbons, the one arising from those bearing hydrogen being easily recognized because it was split by one-bond ^{13}C -H spin-spin coupling

TABLE 1. ^{13}C - ^1H spin-spin couplings (± 0.4 Hz) for **3a**, **3b**, **4a**, **4b** and model compounds **1a**, **1b**, **2a**, **2b**, **5a**, **5b***

	<i>C</i> -1,1'	<i>C</i> -2,2',6,6'	<i>C</i> -3,3',5,5'	<i>C</i> -4,4'			<i>C</i> -2,4,6	<i>C</i> -1,3,5						
2a	³ <i>J</i> 5.3t	ma	¹ <i>J</i> 160.0d ³ <i>J</i> 4.7d	² <i>J</i> 3.5t		1a	¹ <i>J</i> 160.5d ³ <i>J</i> 4.6t	<i>m</i> , <i>w</i> _{1/2} ≈ 9						
2b	³ <i>J</i> 5.5t	mb	¹ <i>J</i> 168.3d ³ <i>J</i> 5.0d	² <i>J</i> 5.0t		1b	¹ <i>J</i> 167.6d ³ <i>J</i> 4.9t	² <i>J</i> ≈ 4.0t ⁴ <i>J</i> ≈ 3.0d						
						5a	<i>C</i> -4' † ¹ <i>J</i> 161.9d ³ <i>J</i> 4 ± 1t	<i>C</i> -3',5' †	<i>C</i> -2',6' †	<i>C</i> -1' †	<i>C</i> -1 †	<i>C</i> -2,6 †	<i>C</i> -3,5 ¹ <i>J</i> 161.6d ³ <i>J</i> 5.3d	<i>C</i> -4 †
						5b	¹ <i>J</i> 167.9d ³ <i>J</i> 5.3t	mb	¹ <i>J</i> 165.9d ³ <i>J</i> 4.7t	² <i>J</i> 4.4t	³ <i>J</i> 7.4t	mb	¹ <i>J</i> 168.1d ³ <i>J</i> 5.6d	² <i>J</i> 5.3t
	<i>C</i> -1'	<i>C</i> -2',6'	<i>C</i> -3',5'	<i>C</i> -4'	<i>C</i> -1	<i>C</i> -2,6	<i>C</i> -3,5	<i>C</i> -4	<i>C</i> -1''	<i>C</i> -2'',6''	<i>C</i> -3'',5''	<i>C</i> -4''		
3a	³ <i>J</i> 5.0t	ma	¹ <i>J</i> 160.1d ³ <i>J</i> 5.0d	² <i>J</i> 3.7t	³ <i>J</i> 5.0t	ma	¹ <i>J</i> 161.4d ³ <i>J</i> 4.0d	² <i>J</i> 3.8t	³ <i>J</i> 7.3t	ma	¹ <i>J</i> 161.1d ³ <i>J</i> 5.1d	² <i>J</i> 3.8t		
3b	³ <i>J</i> 5.0t	mb	¹ <i>J</i> 168.7d ³ <i>J</i> 5.0d	² <i>J</i> 5.0t	³ <i>J</i> 5.0t	mb	¹ <i>J</i> 166.2d ³ <i>J</i> 5.0d	² <i>J</i> 4.7t	³ <i>J</i> 7.3t	mb	¹ <i>J</i> 168.1d ³ <i>J</i> 5.6d	² <i>J</i> 5.3t		
	<i>C</i> -1'	<i>C</i> -2',6'	<i>C</i> -3',5'	<i>C</i> -4'	<i>C</i> -1	<i>C</i> -2,6	<i>C</i> -3,5	<i>C</i> -4	<i>C</i> -1'''	<i>C</i> -2''',6'''	<i>C</i> -3''',5'''	<i>C</i> -4'''		
4a	³ <i>J</i> 4.9t	ma	¹ <i>J</i> 160.3d ³ <i>J</i> 4.7d	² <i>J</i> 3.5t	³ <i>J</i> 5.1t	ma	¹ <i>J</i> 161.9d ³ <i>J</i> 4.6d	² <i>J</i> 4.1t	³ <i>J</i> 7.1t	ma	¹ <i>J</i> 160.7d ³ <i>J</i> 5.0d	² <i>J</i> 4.3t		
4b	³ <i>J</i> 5.5t	mb	¹ <i>J</i> 168.3d ³ <i>J</i> 5.0d	² <i>J</i> 4.8t	³ <i>J</i> 5 ± 1t	mb	¹ <i>J</i> 166.2d ³ <i>J</i> 5 ± 1d	² <i>J</i> 5 ± 1d	³ <i>J</i> 7.5t	mb	¹ <i>J</i> 168.4d ³ <i>J</i> 5.0d	² <i>J</i> 5.5t		
									<i>C</i> -1''	<i>C</i> -2''	<i>C</i> -3''	<i>C</i> -4''	<i>C</i> -5''	<i>C</i> -6''
								4a	³ <i>J</i> 7.2t	² <i>J</i> or ⁴ <i>J</i> 3.8d	¹ <i>J</i> 161.6d ³ <i>J</i> 5.3d	² <i>J</i> 4.3t	¹ <i>J</i> 161.2d ³ <i>J</i> 5.6d	² <i>J</i> or ⁴ <i>J</i> 3.7d
							4b	³ <i>J</i> 7.2t	m		¹ <i>J</i> 170 ± 2d ³ <i>J</i> 5.5d	² <i>J</i> 5.2t	¹ <i>J</i> 168.3d ³ <i>J</i> 5.0d	m

*For the carbons of acetate groups, spin-spin coupling constants (Hz) were: $^{13}\text{CH}_3$, $^1J_{\text{CH}}$ 130.4q; $^{13}\text{C}=\text{O}$, 2J 6.9q, ma: characteristic multiplet (doublet of doublets) due to coupling to one *ortho* and one *para* proton in h.r. spectra of hydroxy compounds; mb: similar multiplet for acetylated compounds; m: multiplet; d: doublet; t: triplet; q: quartet; $w_{1/2}$ line width at half height.
†Poor S/N due to insufficient material for h.r. spectrum.

($^1J_{CH}$) in the h.r. spectra, and occurred at higher field than the signal due to the ring carbons bonded to oxygen, in accord with known ^{13}C substituent effects (17). Similarly, four resonances were observed for the biphenyl ring carbons of **2a** and **2b**, the two at highest field arising from quaternary carbons (C-1, C-1') and carbons carrying hydrogens (C-3, C-3', C-5, C-5'; $^1J_{CH}$). The remaining resonances due to the two types of carbon bonded to oxygen occurred at low field as expected, and could be assigned because the signal for C-2, C-2', C-6, and C-6' was about twice the intensity of the one arising from C-4 and C-4', the relative intensities of the two sets of resonances being independent of the conditions used to record the spectra.

Compound **5b** was synthesized in 5% yield following Glombitza *et al.* (14). Its ^{13}C spectrum contained three signals arising from aromatic carbons bearing hydrogen. Two which appeared as doublets ($^1J_{CH}$) of triplets ($^3J_{CH}$ to two hydrogens) could be assigned to C-4' (δ_c 110.5) and C-2', C-6' (δ_c 107.6), the relative intensity of the former resonance being approximately half that of the latter, a result independent of experimental conditions. The signal for C-3, C-5 was a doublet ($^1J_{CH}$) of doublets ($^3J_{CH}$ to one hydrogen) as expected. Resonances at δ_c 167.9 to δ_c 168.5 were readily ascribed to the acetoxyl carbonyl carbons, and one at δ_c 158.1 clearly belonged to C-1' (the corresponding carbon in diphenyl ether resonates at δ_c 157.9 (17)). A low-intensity signal at δ_c 136.5 could be ascribed to the poorly-relaxed carbon C-1, which is shielded by a *para* and two *ortho* acetoxyl substituents. The resonance at δ_c 151.6 could be assigned to C-3', C-5' as their chemical shift should not differ significantly from that of C-1, C-3, C-5 (δ_c 151.2) in **1b**, the *meta* substituent effects of acetoxyl and phenoxy groups being small (17). The remaining

6a R = H

6b R = COCH₃

two resonances at δ_c 146.8 and δ_c 143.7 were assigned to C-4 and C-2, C-6, respectively, the relative intensity of the former signal being approximately half that of the latter under a variety of experimental conditions. Additional

support for this assignment is provided by the spectrum of diphenyl ether (**17**), where C-2, C-6 (δ_c 119.3) is shielded to a greater extent than C-4 (δ_c 123.6). It is reasonable to assume that the corresponding carbons in **5b** would show comparable differences in chemical shift. A similar analysis assigned the resonances in the ^{13}C spectrum of **5a**, although there was insufficient sample for a h.r. spectrum of the carbons not bonded to hydrogen.

The 18 carbons of **3a** gave rise to 12 resonances in the aromatic region of the 1H broadband-decoupled ^{13}C spectrum. This is consistent with the 1H nmr results discussed above which showed that **3a** was symmetrical, and indirectly but unequivocally established the presence of at least six pairs of chemically equivalent aromatic carbons. Twelve resonances for the aromatic carbons were also observed in the high resolution ^{13}C spectrum of **3b**, but the 1H broadband-decoupled ^{13}C spectrum contained only ten resonances due to superposition of signals (C-1 hidden by C-3', C-5' and C-1' by C-3'', C-5'').

It was apparent from the ^{13}C chemical shift and spin-spin coupling data (Table 1) that **3a** (**3b**) possessed three sets of aromatic carbons. Each set contained two chemically non-equivalent, plus two pairs of chemically equivalent, carbons and gave rise to four resonances. The ^{13}C data for one set of carbons (**3a**, **3b**; C-1' through C-6') was virtually identical to that for **2a** (**2b**), and that of another set (**3a**, **3b**; C-1 through C-6) differed only slightly, and predictably (see comments on chemical shift calculations below). Similarly the ^{13}C data for the carbons in the remaining set (**3a**, **3b**; C-1'' through C-6'') were almost indistinguishable from those obtained for C-1 through C-6 of **5a** (**5b**).

7a R = H^d7b R = COCH₃^{a,b,d,e}

The ^{13}C chemical shifts expected for C-1 through C-6 of **3a** and **3b** were calculated from the known chemical shifts for **1a**, **1b**, **2a**, **2b**, **5a**, and **5b** as follows. Comparison of **1a**, **1b** and **5a**, **5b** yields the changes in chemical shift at all

the carbons of **1a,1b** due to replacement of one hydroxy (or acetoxy) substituent by a 2,4,6-trihydroxy (or triacetoxy) phenoxy group: at the directly bonded carbon $+2.2(+6.9)$ ppm; *ortho* $-1.0(-5.2)$ ppm; *meta* $0.0(+0.4)$ ppm; *para* $+1.6(-2.3)$ ppm. These chemical shift changes applied to **2a,2b** for the same change of substitution at C-4 predict the following chemical shifts for C-1 through C-6 of **3a,3b**. The close correspondence of measured and predicted (in parentheses) shifts provides further support for the ^{13}C resonance assignments and the structures: **3a** C-1 104.2 (104.2), C-2,C-6 159.4 (159.2), C-3,C-5 97.4 (97.2), C-4 162.5 (162.3); **3b** C-1 113.9 (113.5), C-2,C-6 150.0 (149.8), C-3,C-5 108.8 (108.6), C-4 157.7 (157.5).

In addition, hydrogen to deuterium exchange in D_2O eliminated ^{13}C - ^1H spin-spin coupling from all resonances except those assigned to C-1 through C-6 of **3a**, which partially retained coupling. This confirms that the latter carbons belonged to the same ring. The centre ring of **3a** would be expected to exchange hydrogen less rapidly, as it provides less opportunity for keto-enol tautomerism. The combined evidence therefore establishes that **3a** is 4-(2'',4'',6''-trihydroxyphenoxy)-2,2',4',6,6',-pentahydroxybiphenyl.

The ^1H nmr spectrum for **4a** (solvent acetone- d_6) contained three singlet resonances at δ 6.01, 6.03, and 6.14 due to three pairs of chemically equivalent aromatic hydrogens. Two of these (δ 6.01, 6.03) corresponded closely to the signals for hydrogens at C-3, C-3', C-5, C-5' of **2a** and C-3, C-3', C-5, C-5', C-3'', C-5'' of **3a**, while the third (δ 6.14) was similar apart from a small downfield shift. The spectrum also contained an AB multiplet for two *meta*-coupled aromatic hydrogens (centroid δ 6.01, $\Delta\nu$ 32.7 Hz, $^3J_{\text{HH}}$ 2.8 Hz) and a broad OH signal (δ 6.88) integrating for 10 protons.

The ^{13}C spectrum of **4a** contained 18 resonances, 12 of which corresponded almost exactly in chemical shift, relative intensity (recorded under widely varying conditions), and ^{13}C -H coupling (Table 1) with the resonances for **3a**. This information combined with the known molecular formula established that **4a** contained 2,2',4',6,6'-pentahydroxybiphenoxy and 2,4,6-trihydroxyphenoxy residues, which were asymmetrically substituted on another aromatic ring containing $\text{C}_6\text{H}_4\text{O}_3$. Similarly the ^{13}C spectrum of **4b** supported these conclusions.

The presence of two *meta*-coupled hydrogens limits the choice of structure of the acetylated material to five possibilities, **4b**, and **4d** to **4g**. The choice of **4b** as the correct structure was based on a comparison of the measured ^{13}C chemical shifts for the carbons of the asymmetrically substituted aromatic ring (C-1'' to C-6'') with predicted shifts derived from measurements of acetylated model compounds, as follows. Firstly, three independent predictions were made of the ^{13}C shifts for **4h**.

(a) The average effect of replacement of an acetoxy substituent on a ring by a 2,4,6-triacetoxyphenoxy group was estimated by comparing the measured δ_c values for **1b, 5b; 2b, 3b; 1b, 8** (18); **5b, 8** (18): at substituent $+6.9 \pm 0.2$ ppm; *ortho* -5.4 ± 0.4 ppm; *meta* $+0.4 \pm 0.2$ ppm; *para* -2.6 ± 0.7 ppm. This information was then used to predict the ^{13}C chemical shifts for **4h** by applying the substituent effects above, in reverse, to allow for substitution of an acetoxy group at C-1' of **7b**.

(b) The measured shifts of **5b** and **7b** were compared to estimate the substituent effects of acetoxy-substitution at C-4' of **5b** and these were applied to predict the result of substitution of an acetoxy group at C-2 of **1b**.

(c) **1b** and **9** were compared to obtain the effects of acetoxy substitution at C-5 of **9**, and these were used to predict the effect of acetoxy substitution at C-5 of **10**. The averages of these predictions (a,b,c) of the δ_c values for **4h** were: C-1'' 132.2 ± 0.9 ; C-2'',6'' 143.3 ± 0.6 ; C-3'',5'' 114.4 ± 0.5 ; C-4'' 147.5 ± 0.6 ppm. When these values were compared with the measured δ_c values for C-1'' to C-6'' of **3b**, the effect of replacing an acetoxy substituent by a (2,2',4',6,6')-pentaacetoxy biphenoxy group was obtainable: at substituent $+4.4(\pm 0.9)$; *ortho* $+0.5(\pm 0.6)$; *meta* $+0.8(\pm 0.5)$; *para* $-0.6(\pm 0.6)$ ppm. The predicted δ_c for **4h** and the substituent effects above were used to predict the chemical shifts for the five alternative structures **4b, 4d** to **4g**. For example, δ_c for C-1'' of **4b** is given by δ_c for C-1'' of **4h** (132.2 ppm), plus the effect of replacing the acetoxy substituent at C-1'' with a 2,2',4',6,6'-pentaacetoxy biphenoxy group ($+4.4$ ppm) and the effect of replacing the *ortho*-acetoxy at C-2'' by a 2,4,6-triacetoxyphenoxy substituent (-5.4 ppm). The total, 131.2 ppm, is the predicted δ_c . The nearest experimental δ_c values for this ring are compared with predictions in Table 2, which also shows the sum of

TABLE 2. Comparison of experimental δ_c (ppm) for C-1'' through C-6'' of **4b**, **4d-g** with nearest predicted values

Experimental δ_c (ppm)	Nearest predicted δ_c for alternative structures (ppm)				
	4b	4d	4e	4f	4g
112.2	109.8 C-3''	109.5 C-5''	108.4 C-3''	114.2 C-3''	108.4 C-3''
113.8	112.6 C-5''	112.3 C-3''	112.3 C-5''	115.3 C-5''	109.5 C-5''
133.4	131.2 C-1''	126.2 C-1''	127.3 C-1''	138.7 C-2''	130.1 C-1''
144.1	144.2 C-6''	144.5 C-2''	148.1 C-6''	139.6 C-1''	144.5 C-2''
147.3	147.3 C-4''	151.0 C-6''	148.7 C-4''	142.3 C-6''	148.1 C-6''
150.7	150.7 C-2''	152.3 C-4''	151.0 C-2''	145.7 C-4''	155.2 C-4''
$\Sigma (\text{Exptl.} - \text{predicted}) $	5.9	17.1	17.1	23.3	17.3

TABLE 3. Predicted δ_c for C-1'' to C-6'' of **4a** and **4b** (ppm)

		C-1''	C-2''	C-3''	C-4''	C-5''	C-6''
4a	Predicted	125.7	155.2	98.0	156.9	100.5	153.0
	Measured	126.7	154.9	96.7	156.7	100.5	153.3
4b	Predicted	131.2	150.7	109.8	147.3	112.6	144.2
	Measured	133.4	150.7	112.2	147.3	113.8	144.1

the absolute magnitudes of the differences between predicted and experimental values. Thus, structure **4b** is clearly favoured. The predicted shifts for **4b** (or **4a**) may also be calculated from the experimental shifts for **3b** (**3a**) and the substituent effects under (a) above (or the corresponding substituent effect for replacement of OH by a 2,4,6-trihydroxyphenoxy substituent, estimated from experimental data for **1a**, **5a**; **2a**, **3a**), again giving close agreement (Table 3) with the other predicted shifts (Table 2, structure **4b**) and with the experimentally measured values.

F. vesiculosus from Nova Scotia has been shown to contain 4-(2'',4'',6''-trihydroxyphenoxy)-2,2',4',6,6'-pentahydroxybiphenyl **3a** and

4-(2''-(2''',4''',6'''-trihydroxyphenoxy)-4'',6''-dihydroxyphenoxy)-2,2',4',6,6'-pentahydroxybiphenyl **4a** in addition to compounds **1a** and **2a** (6). If a terphenyl or quaterphenyls (7) were present in our extracts, their trimethylsilyl ethers should appear at *m/e* 1022 and 1362, respectively. In fact, only a trace of an otherwise unidentified compound appeared at *m/e* 1022, and no *m/e* 1362 ion was observed following gc/ms (Ragan and Craigie, unpublished) and high resolution ms with photoplate ion beam integration. Similarly, the ether linked C₁₂ and C₁₈ polyphloroglucinols reported from *Bifurcaria bifurcata* (9, 10), *Cystoseira tamariscifolia* (14), *Halidrys siliquosa* (11), and *Laminaria ochroleuca* (18) must occur only in minute quantities, if at all, in our *F. vesiculosus*.

Acknowledgments

We thank Dr. W. D. Jamieson and Mr. D. Embree for the high resolution mass measurements and for helpful discussions. M.A.R. acknowledges financial support for his graduate studies from Dalhousie University and the National Research Council of Canada.

1. E. CRATO. Ber. Dtsch. Bot. Ges. **10**, 295 (1892).
2. M. A. RAGAN. Bot. Mar. **19**, 145 (1976).
3. C. OGINO and Y. TAKI. J. Tokyo Univ. Fish. **43**, 1 (1957).
4. J. S. CRAIGIE and J. McLACHLAN. Can. J. Bot. **42**, 23 (1964).

5. K.-W. GLOMBITZA, H.-U. RÖSENER, H. VILTER, and W. RAUWALD. *Planta Med.* **24**, 301 (1973).
6. M. A. RAGAN and J. S. CRAIGIE. *Can. J. Biochem.* **54**, 66 (1976).
7. K.-W. GLOMBITZA, H.-W. RAUWALD, and G. ECKHARDT. *Phytochemistry*, **14**, 1403 (1975).
8. K.-W. GLOMBITZA. VIIIth International Seaweed Symposium, Bangor, Wales. 1974.
9. K.-W. GLOMBITZA and H.-U. RÖSENER. *Phytochemistry*, **13**, 1245 (1974).
10. K.-W. GLOMBITZA, H.-U. RÖSENER, and M. KOCH. *Phytochemistry*, **15**, 1279 (1976).
11. K.-W. GLOMBITZA and E. SATTLER. *Tetrahedron Lett.* **43**, 4277 (1973).
12. J. B. GRUTZNER and R. E. SANTINI. *J. Magn. Reson.* **19**, 173 (1975).
13. G. C. LEVY and G. L. NELSON. Carbon-13 nuclear magnetic resonance for organic chemists. Wiley-Interscience, New York, NY. 1972.
14. K.-W. GLOMBITZA, H.-U. RÖSENER, and D. MÜLLER. *Phytochemistry*, **14**, 1115 (1975).
15. M. A. RAGAN. Physodes and the phenolic compounds of brown algae. Ph.D. thesis, Dalhousie University, Halifax, N.S. 1976.
16. L. M. JACKMAN and S. STERNHELL. Applications of nuclear magnetic resonance spectroscopy in organic chemistry. 2nd ed. Pergamon, New York, NY. 1969.
17. J. B. STOTHERS. Carbon-13 NMR spectroscopy. Academic Press, New York, NY. 1972.
18. K.-W. GLOMBITZA, M. KOCH, and G. ECKHARDT. *Phytochemistry*, **15**, 1082 (1976).
19. E. SATTLER. Thesis, Rheinischen Friedrich-Wilhelms-Universität zu Bonn. 1974.